



(11) **EP 1 473 368 A1**

(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
03.11.2004 Bulletin 2004/45

(21) Application number: 03706925.9

(22) Date of filing: 06.02.2003

(51) Int Cl.7: **C12N 15/53, C12N 9/02,**  
**C12N 5/10, C12N 1/15,**  
**C12N 1/19, C12N 1/21,**  
**C12P 7/40**

(86) International application number:  
**PCT/JP2003/001240**

(87) International publication number:  
**WO 2003/066863 (14.08.2003 Gazette 2003/33)**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR**  
**HU IE IT LI LU MC NL PT SE SI SK TR**  
Designated Extension States:  
**AL LT LV MK RO**

(30) Priority: 06.02.2002 JP 2002030127  
26.09.2002 JP 2002281236

(71) Applicant: **SHOWA DENKO K.K.**  
**Tokyo 105-8518 (JP)**

(72) Inventors:  
• **KAMACHI, Harumi, Corporate R & D Center**  
**Chiba-shi, Chiba 267-0056 (JP)**  
• **ESAKI, Nobuyoshi**  
**Otsu-shi, Shiga 520-2279 (JP)**  
• **KURIHARA, Tatsuo,**  
**Fushimi Godo Shukusha 533**  
**Kyoto-shi, Kyoto 612-8104 (JP)**

(74) Representative: **Strehl Schübel-Hopf & Partner**  
**Maximilianstrasse 54**  
**80538 München (DE)**

(54) **ALPHA-SUBSTITUTED-ALPHA, BETA-UNSATURATED CARBONYL COMPOUND**  
**REDUCTASE GENE**

(57) The present invention relates to: a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound which contains a DNA sequence encoding an amino acid sequence represented by SEQ ID NO: 20 and an amino acid sequence represented by SEQ ID NO: 21; an enzyme which is a product of the gene; a plasmid and a transformant each containing the gene DNA; and a method of reducing an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound using the transform-

ant. According to claim the present invention, there is provided an enzyme gene which is useful in producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound optically active at the  $\alpha$ -position by hydrogenating an  $\alpha,\beta$ -carbon double bond of an  $\alpha$ -substituted carbonyl compound, which is a compound prochiral at the alpha-position, and an enzyme which is a gene product thereof.

**EP 1 473 368 A1**



(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
03.11.2004 Bulletin 2004/45

(21) Application number: 03706925.9

(22) Date of filing: 06.02.2003

(51) Int Cl.<sup>7</sup>: **C12N 15/53**, **C12N 9/02**,  
**C12N 5/10**, **C12N 1/15**,  
**C12N 1/19**, **C12N 1/21**,  
**C12P 7/40**

(86) International application number:  
**PCT/JP2003/001240**

(87) International publication number:  
**WO 2003/066863 (14.08.2003 Gazette 2003/33)**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR**  
**HU IE IT LI LU MC NL PT SE SI SK TR**  
Designated Extension States:  
**AL LT LV MK RO**

(30) Priority: **06.02.2002 JP 2002030127**  
**26.09.2002 JP 2002281236**

(71) Applicant: **SHOWA DENKO K.K.**  
**Tokyo 105-8518 (JP)**

(72) Inventors:  
• **KAMACHI, Harumi, Corporate R & D Center**  
**Chiba-shi, Chiba 267-0056 (JP)**  
• **ESAKI, Nobuyoshi**  
**Otsu-shi, Shiga 520-2279 (JP)**  
• **KURIHARA, Tatsuo,**  
**Fushimi Godo Shukusha 533**  
**Kyoto-shi, Kyoto 612-8104 (JP)**

(74) Representative: **Strehl Schübel-Hopf & Partner**  
**Maximilianstrasse 54**  
**80538 München (DE)**

(54) **ALPHA-SUBSTITUTED-ALPHA, BETA-UNSATURATED CARBONYL COMPOUND**  
**REDUCTASE GENE**

(57) The present invention relates to: a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound which contains a DNA sequence encoding an amino acid sequence represented by SEQ ID NO: 20 and an amino acid sequence represented by SEQ ID NO: 21; an enzyme which is a product of the gene; a plasmid and a transformant each containing the gene DNA; and a method of reducing an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound using the transform-

ant. According to claim the present invention, there is provided an enzyme gene which is useful in producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound optically active at the  $\alpha$ -position by hydrogenating an  $\alpha,\beta$ -carbon double bond of an  $\alpha$ -substituted carbonyl compound, which is a compound prochiral at the alpha-position, and an enzyme which is a gene product thereof.

## Description

## TECHNICAL FIELD

5 [0001] The present invention relates to a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, and enzymes as gene products thereof. More specifically, the present invention relates to a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound having activities of producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound by hydrogenating an  $\alpha$ ,  $\beta$ -carbon-carbon double bond of an  $\alpha$ -substituted carbonyl compound characterized in that the gene is derived from at least one microorganism selected from the group consisting of the genus Acetobacter, Actinomyces, Acinetobacter, Agrobacterium, Aeromonas, Alcaligenes, Arthrobacter, Azotobacter, Bacillus, Brevibacterium, Burkholderia, Cellulomonas, Corynebacterium, Enterobacter, Enterococcus, Escherichia, Flavobacterium, Gluconobacter, Halobacterium, Halococcus, Klebsiella, Lactobacillus, Microbacterium, Micrococcus, Micropolyspora, Mycobacterium, Nocardia, Pseudomonas, Pseudonocardia, Rhodococcus, Rhodobacter, Serratia, Staphylococcus, Streptococcus, Streptomyces, and Xanthomonas, and enzymes as gene products thereof.

15 [0002] Further, the present invention relates to a reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound derived from genus Pseudomonas or Burkholderia microorganisms, in particular Pseudomonas sp. SD810 strain, Pseudomonas sp. SD811 strain, Pseudomonas sp. SD812 strain, and Burkholderia sp. SD816 strain, or enzymes as gene products thereof having the above activities.

20 [0003] Furthermore, the present invention relates to a reductase gene and an enzyme as a gene product thereof, which are useful in producing an  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound optically active at the  $\alpha$ -position by stereoselectively hydrogenating a carbon-carbon double bond in a corresponding  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond, which is a molecule prochiral at the  $\alpha$ -position.

25 [0004] The novel enzyme gene and the enzyme as the product thereof can be used in the field of production of optically active carbonyl compounds including various optically active saturated carboxylic acids (having the S- or R-form absolute configurations at their respective  $\alpha$ -positions with substituted groups, respectively) or amides. The optically active carbonyl compounds are highly valuable chiral building blocks, which can be hardly prepared by classical chemical processes, and in particular the compounds are useful materials as raw materials of medical and agricultural chemicals.

## 30 BACKGROUND ART

[0005] In recent years, much attention has been paid on the method of producing various compounds, particularly optically active substances, by the microbial reduction of carbon-carbon double bonds. To this end, many publications have reported various processes of producing a corresponding  $\alpha,\beta$ -saturated carbonyl compound having a substituent at the  $\alpha$ -position from a carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond and having a substituent at the  $\alpha$ -position by microbially reducing the carbon-carbon double bond (See e.g., Hoppe-Seyler's Z. Physiol. Chem. 362, 33 (1981); Arch. Microbiol. 135, 51 (1983); Helv. Chim. Acta., 62, 455 (1979); J. Ferm. Bioeng., 84, 195 (1997)).

35 [0006] However, no example has been provided with respect to the separation and identification of reductase from active microorganisms used in these processes. Firstly, few studies have been performed on enzymes belonging to this group because of difficulties in separation and identification due to their instability. The enzyme of the present invention has not been an exceptional case, so that the separation and identification of the enzyme has been impossible in the conventional process because of its rapid inactivation.

40 [0007] On this account, such a disadvantage has been difficult to mitigate by a genetic- or metabolic-engineering approach when the reductase is used in the production of a chemical compound. Therefore, an effective improvement in production process has been hardly conducted.

## DISCLOSURE OF THE INVENTION

50 [0008] Therefore, an object of the present invention is to provide a catalytic enzyme useful in producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound from an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond by microbial reduction of the carbon-carbon double bond and to provide a gene of the catalytic enzyme.

55 [0009] Thorough screening from soil has allowed the inventors of the present invention to find that, surprisingly, microorganisms, each of which is capable of producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound from an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond by reduction of the carbon-carbon double bond, are distributed over a relatively wide genus range of the aerobic and facultative anaerobic bacteria (e.g., JP 10-224821 A).

[0010] In particular, it has been found that a large number of strains having the above enzymatic activity are present

in microorganisms belonging to the genera Pseudomonas and Burkholderia, and some of these strains can reduce an  $\alpha$ -halocarbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond to thereby produce an extremely high-purity  $\alpha$ -halo- $\alpha,\beta$ -saturated carbonyl compound having the S absolute configuration at the  $\alpha$ -position.

[0011] Furthermore, the inventors of the present invention have succeeded in establishing a method of producing optically active  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compounds using these active microorganisms and dedicated to studying for identification of reductase itself and also for identification of a gene thereof to improve the production process. As a result, the inventors of the present invention have succeeded in identifying a catalytic enzyme, revealing the action mechanism of the reductase, and collecting microorganisms having high activity, resulting in the completion of the present invention.

[0012] In other words, the present invention relates to a reductase gene, plasmid, transformant, a protein, a method of producing a gene that encodes the protein, and a reductase gene for an  $\alpha,\beta$ -unsaturated carbonyl compound.

1. A gene including: DNA having a base sequence represented by SEQ ID NO: 19 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.

2. A gene including: DNA having a base sequence represented by SEQ ID NO: 17 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.

3. A gene including: DNA having a base sequence represented by SEQ ID NO: 18 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.

4. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, characterized by including a DNA sequence encoding an amino acid sequence represented by SEQ ID NO: 20 and an amino acid sequence represented by SEQ ID NO: 21.

5. A gene that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising an amino acid sequence represented by SEQ ID NO: 20 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids.

6. A gene that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising an amino acid sequence represented by SEQ ID NO: 21 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids.

7. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of 1 to 6, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is derived from at least one microorganism selected from the group consisting of the genus Acetobacter, Actinomyces, Acinetobacter, Agrobacterium, Aeromonas, Alcaligenes, Arthrobacter, Azotobacter, Bacillus, Brevibacterium, Burkholderia, Cellulomonas, Corynebacterium, Enterobacter, Enterococcus, Escherichia, Flavobacterium, Gluconobacter, Halobacterium, Halococcus, Klebsiella, Lactobacillus, Microbacterium, Micrococcus, Micropolyspora, Mycobacterium, Novcardia, Pseudomonas, Pseudonocardia, Rhodococcus, Rhodobacter, Serratia, Staphylococcus, Streptococcus, Streptomyces, and Xanthomonas.

8. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 7, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is derived from a Pseudomonas microorganism.

9. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 7, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is originated from a Burkholderia microorganism.

10. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD810 strain.

11. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD811 strain.


12. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD812 strain.

13. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 9, in which the Burkholderia microorganism is Burkholderia sp. SD816 strain.

14. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of 1 to 13, in which the reductase has a catalytic activity to reduce a carbon-carbon double bond to produce an S-form compound chiral at an  $\alpha$ -position.


15. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of 1 to 14, in which:

the  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is a compound represented by the following general formula (1)



wherein  $\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  each independently represent a hydrogen atom, a halogen atom, a linear or branched aliphatic hydrocarbon group having 1 to 6 carbon atoms, a linear or branched alkoxy group having 1 to 6 carbon atoms, a hydroxyl group, a carboxyl group, an aromatic group or a nitrogen-, oxygen-, or sulfur-containing heterocyclic group which may be substituted, and  $\text{R}^4$  represents a hydroxyl group, a linear or branched alkoxy group having 1 to 3 carbon atoms, or a primary, secondary, or tertiary amino group, provided that  $\text{R}^3$  is not a hydrogen atom; and


a reduced compound is a compound represented by the following general formula (2)



wherein  $\text{R}^1$  to  $\text{R}^4$  have the same meanings as those defined above.


16. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to 15, in which:

the  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is an  $\alpha$ -haloacrylic acid represented by the following general formula (1)



wherein  $\text{R}^1$  and  $\text{R}^2$  represent hydrogen atoms,  $\text{R}^3$  represents a halogen atom, and  $\text{R}^4$  represents a hydroxyl group; and

the reduced compound is an  $\alpha$ -halopropionic acid having an S absolute configuration represented by the following general formula (2)



wherein R<sup>1</sup> to R<sup>4</sup> have the same meanings as those defined above.

17. A reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to 16, in which R<sup>3</sup> represents a bromine atom.

18. A reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to 16, in which R<sup>3</sup> represents a chlorine atom.

19. A reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to 16, in which R<sup>3</sup> represents a fluorine atom.

20. A plasmid, characterized by containing a DNA of a reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to any one of 1 to 19.

21. A plasmid, characterized by containing a reductase gene for an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to any one of 1 to 19 and a gene for an enzyme functioning with an NADPH as a co-enzyme.

22. A transformant transformed with a plasmid according to 20 or 21.

23. A transformant including a product transformed by a plasmid according to 20, and a plasmid containing a gene for an enzyme functioning with an NADPH as a co-enzyme.

24. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound wherein the protein is an expression product of a reductase gene for the  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound according to any one of 1 to 19, or a protein having deletion, substitution, or addition of one or more amino acids thereof and having a reduction activity to an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound.

25. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound containing an amino acid sequence represented by SEQ ID NO: 20 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids in the said amino acid sequence.

26. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound containing an amino acid sequence represented by SEQ ID NO: 21 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids in the said amino acid sequence.

27. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 631 and a base sequence selected from base sequences located downstream of a base at position 3, 543 in the base sequence represented by SEQ ID NO: 19, where both the base sequences extend in opposite directions to each other.

28. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 631 and a base sequence selected from base sequences located downstream of a base at position 2,274 in base sequences represented by SEQ ID NO: 19, where both base sequences extend in opposite directions to each other.

29. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 2,547 and a base sequence selected from base sequences located downstream of a base at position 3,543 in base sequences represented by SEQ ID NO: 19, where both base sequences extend in opposite directions so as to be reversed strands with respect to each other.

30. A method of reducing an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -unsaturated carbonyl compound, comprising:

using a culture and/or treated product of a transformant according to 22 or 23.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0013]

- 5 Fig. 1 is a graph showing different reaction curves with different carbon sources in the cultures of Pseudomonas sp. SD811 strain;  
 Fig. 2 is a graph showing different reaction curves with different carbon sources in the cultures of Burkholderia sp. SD816 strain;  
 Fig. 3 is a two-dimensional electrophoretic image that shows the results in the comparison among proteins from  
 10 cultures of Burkholderia sp. SD816 strain with different carbon sources;  
 Fig. 4 is a schematic diagram that represents a positional relationship between a CAA43 gene and a CAA67 gene and also represents the positions of the respective DNA fragments (the number of each fragment represents SEQ ID NO); and  
 Fig. 5 is a schematic diagram that represents the construction of an expression vector for a reductase gene responsible for asymmetric reduction.

## DETAILED DESCRIPTION OF THE INVENTION

20 [0014] A reductase gene for an  $\alpha$ -substituted-  $\alpha,\beta$ -unsaturated carbonyl compound and enzymes as products thereof according to the present invention exist in microorganisms belonging to any one of the genus Acetobacter, Actinomyces, Acinetobacter, Agrobacterium, Aeromonas, Alcaligenes, Arthrobacter, Azotobacter, Bacillus, Brevibacterium, Burkholderia, Cellulomonas, Corynebacterium, Enterobacter, Enterococcus, Escherichia, Flavobacterium, Gluconobacter, Halobacterium, Halococcus, Klebsiella, Lactobacillus, Microbacterium, Micrococcus, Micropolyspora, Mycobacterium, Nocardia, Pseudomonas, Pseudonocardia, Rhodococcus, Rhodobacter, Serratia, Staphylococcus, Streptococcus, Streptomyces, and Xanthomonas.

25 [0015] Preferably they are derived from Pseudomonas or Burkholderia microorganisms.

[0016] Original microorganisms used in the present invention may be any strains as far as they have the activity of reducing an  $\alpha,\beta$ -carbon-carbon double bond of an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond. Preferable examples of the microorganisms include, but not particularly limited to, Pseudomonas sp. SD810 strain, Pseudomonas sp. SD811 strain, Pseudomonas sp. SD812 strain, and Burkholderia sp. SD816 strain.

30 [0017] Among those, Pseudomonas sp. SD811 strain or Burkholderia sp. SD816 strain is particularly preferably used in terms of comparatively high reduction activity.

[0018] The microorganisms used, such as Pseudomonas sp. SD810 strain, Pseudomonas sp. SD811 strain, Pseudomonas sp. SD812 strain, and Burkholderia sp. SD816 strain, which are isolated from soil, have their own activities of decomposing and assimilating various carbonyl compounds.


35 [0019] These microorganisms, Pseudomonas sp. SD810 strain, Pseudomonas sp. SD811 strain, and Pseudomonas sp. SD812 strain, were deposited with the National Institute of Bioscience and Human-Technology under the accession numbers BP-6767 (FERMBP-6767) (transferred from accession number 16746 (FERM-16746)), BP-6768 (FERM BP-6768) (transferred from accession number 16747 (FERM-16747)), and BP-6769 (FERM BP-6769) (transferred from accession number 16748 (FERM-BP-6769)), respectively. In addition, Burkholderia sp. SD816 strain is deposited with the National Institute of Bioscience and Human-Technology under the accession number BP-6770 (FERM BP-6770).

40 [0020] Those strains may be isolated and cultured by the conventional procedures including those specifically described in JP 10-224821 A.

[0021] The active microorganisms described above may show variations in their reduction activities depending on their culture conditions. That is, each of the microorganisms shows different activities on the reduction of an  $\alpha,\beta$ -carbon-carbon double bond between the case where the microorganism is cultured using an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond (i.e., reduction substrate) as a carbon source and the case where the microorganism is cultured using a typical carbon source such as a saccharide. That is, a microorganism cultured using a reduction substrate as a carbon source may show a high reduction activity from the beginning of the reaction. It suggests that the reductase is induced partly or wholly with the reduction substrate, so that an analysis on such a difference will lead to the identification of the reductase.

50 [0022] A carbon source in a culture medium for obtaining a microorganism having a high reduction activity may be a compound represented by the general formula (1).

55



[0023] In the formula (1), R<sup>1</sup> and R<sup>2</sup> each independently represent a hydrogen atom, a halogen atom, a linear or branched aliphatic hydrocarbon group having 1 to 6 carbon atoms, a linear or branched alkoxy group having 1 to 6 carbon atoms, a hydroxyl group, a carboxyl group, an aromatic group or a saturated or unsaturated nitrogen-, oxygen-, or sulfur-containing heterocyclic group which may be substituted. Preferably R<sup>1</sup> and R<sup>2</sup> are hydrogen atoms;

[0024] R<sup>3</sup> represents a halogen atom, a linear or branched aliphatic hydrocarbon group having 1 to 6 carbon atoms, a linear or branched alkoxy group having 1 to 6 carbon atoms, a hydroxyl group, a carboxyl group, an aromatic group or a saturated or unsaturated nitrogen-, oxygen-, or sulfur-containing heterocyclic group which may be substituted, preferably a halogen atom, in particular a chlorine atom or a bromine atom.

[0025] R<sup>4</sup> represents a hydroxyl group, a linear or branched alkoxy group having 1 to 4 carbon atoms, or a primary, secondary, or tertiary amino group, preferably a hydroxyl group.

[0026] Specific examples of the compound include  $\alpha$ -chloroacrylic acid,  $\alpha$ -bromoacrylic acid, 2-chloro-2-butenic acid, 2-bromo-2-butenic acid, 2-chloro-2-pentenoic acid, 2-bromo-2-pentenoic acid, and methyl esters and ethyl esters thereof. Of these,  $\alpha$ -chloroacrylic acid and  $\alpha$ -bromoacrylic acid are preferred.

[0027] More specifically, bacterial cells having a high reduction activity can be obtained by: inoculating a strain in 5 ml of minimal medium prepared by adding 2 g/l of an  $\alpha,\beta$ -unsaturated carbonyl compound having a substituent at the  $\alpha$ -position, such as  $\alpha$ -chloroacrylic acid, as a substantially only carbon source to an inorganic salt culture medium (e. g., (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>: 2 g/l, NaH<sub>2</sub>PO<sub>4</sub>: 1 g/l, K<sub>2</sub>HPO<sub>4</sub>: 1 g/l, MgSO<sub>4</sub>: 0.1 g/l, yeast extract: 0.5 g/l) used for normal bacteria; and incubating the bacteria at 28°C for 12 to 72 hours while shaking. On the other hand, when the bacteria cells are incubated such that only the carbon source in the above culture conditions is replaced with a metabolic product of the reduction substrate, for example lactic acid when the carbon source is a substituted acrylic acid such as  $\alpha$ -chloroacrylic acid, bacterial cells having no reduction activity can be obtained at the beginning of the reaction.

[0028] These bacterial cells are collected by centrifugation and disrupted by the conventional method such as French press to obtain a cell-free extract. Then, the cell-free extract is subjected to column chromatography to make a comparison between the migration patterns of separated proteins, exhibiting different proteins between the bacterial cells incubated under different conditions.

[0029] Among the proteins produced from the bacterial cells incubated using the reduction substrate, proteins having increased amounts of production may be isolated and then the activity thereof may be measured, allowing the identification of the desired enzymes. In general, however, such enzymes show low stability in the state of a cell-free extract. Therefore, the activities of the enzymes disappear comparatively quickly, so that the separation and identification of the enzymes will be difficult in many cases. This fact is one of the causes involved in stagnation in research on enzymes belonging to the group of the above enzymes compared with other stable enzymes.

[0030] In this case, the activity of the enzyme may be retained by carrying out the isolation procedures under nitrogen atmosphere. Alternatively, however, there is an effective process in which a partial sequence of a gene is revealed, a target gene is cloned using a DNA base sequence estimated from the partial sequence as a probe, and the gene is then expressed to obtain a significant amount of the protein, followed by analyzing the protein for its activity or the like.

[0031] In other words, the production patterns of proteins separated from cell-free extracts using different carbon sources by two-dimensional protein electrophoresis or the like are compared and then a protein being increased in bacterial cells incubated with a reduction substrate is found. Subsequently, the protein thus obtained is transferred to a PVDF membrane or the like, followed by analyzing the N-terminal sequence of the protein using a vapor-phase Edman degradation apparatus or the like. A DNA base sequence is estimated from the resulting N-terminal sequence and the corresponding oligonucleotide is then synthesized to prepare a probe useful for acquiring genes for a group of reductase enzymes from chromosomes (i.e., a DNA fragment labeled with an identifiable marker, which can be used for finding out DNA having a specific sequence).

[0032] The reductase gene of the present invention can be easily obtained by the conventional methods such as Southern hybridization generally used in genetic engineering using a DNA probe prepared as described above. More specifically, DNA extracted from the above microorganism (including plasmid if the DNA exists in chromosome and in



plasmid) is cut into fragments by appropriate restriction enzymes. The resulting fragments are separated in size by means of agarose gel electrophoresis or the like and then transferred on a nitrocellulose membrane, followed by subjecting the transferred fragments to hybridization with a probe labeled with an identifiable marker (here, the term "hybridization" means the formation of a double strand DNA when there is high base complementarity between DNA sequences, and is also referred to as "pairing"), resulting in a fragment that hybridizes the probe in a specific manner, or a DNA fragment that contains a target gene. In this case, although the gene may be cut into partial fragments, the entire gene can be obtained by employing the same detection method with different kinds of restriction enzymes, using a previously obtained fragment as a probe, or the like.

[0033] If a hybridization method is applied on genes for a group of reductases of the present invention, although appropriate conditions may be different depending on the length of DNA to be hybridized, a sufficiently specific hybridization result will be obtained under stringent conditions of about 40°C to 70°C, preferably 47°C to 60°C within a salt concentration range of a typical hybridization solution.

[0034] The genes for a group of reductases of the present invention can be also obtained easily by forming primers that hybridize on appropriate sites of the genes and peripheral sequences of the genes; and performing a polymerization chain reaction (PCR) using the microbial DNA as a template.

[0035] The term "primer" used herein is a fragment that is hybridized on a target DNA sequence to be replicated and functions as the initiation point of DNA synthesis. A primer is indispensable in initiation of DNA replication because enzymatic DNA synthesis proceeds such that DNA polymerase catalyses the diester-binding of deoxyribonucleotide on the 3'-OH position of the primer hybridized on the template DNA. A primer is used even for a polymerase chain reaction (PCR), where efficient replication of the target DNA depends on the selection of such a primer.

[0036] A primer, which can be used in the present invention, is not limited to specific one as far as it will be hybridized on the reductase gene of the present invention and the peripheral sequence of the gene and will function as the initiation point for DNA synthesis. For example, there are no limitations on the degree of the sequence complementarity of the fragment, the length of the fragment, modifications to the fragment, and the like. For any purpose, for example, a primer that contains an adaptor sequence for connecting a fragment generated to a plasmid, a primer modified by a fluorescent substance for facilitating the detection of a gene fragment generated, or the like can be designed and used at will.

[0037] A pair of primers useful for obtaining genes for a group of reductases in the present invention is a combination of one having a base sequence containing a sequence upstream of the base at position 631, which is a first base of the initiation codon of the upstream gene among the base sequences represented in SEQ ID NO: 19, and the other having a base sequence downstream of the base at position 3, 543, which is a third base of the termination codon of the downstream gene, such that the primer strands extend in opposite directions to each other. Another pair of primers useful for obtaining genes for a group of reductases is a combination of one having a base sequence containing a sequence upstream of the base at position 631, which is a first base of the initiation codon of the upstream gene among the base sequences represented in SEQ ID NO: 19, and the other having a base sequence downstream of the base at position 2,274, which is a third base of the termination codon of the upstream gene, such that the primer strands extend in opposite directions to each other. Further another pair of primers useful for obtaining genes for a group of reductases is a combination of one having a base sequence containing a sequence upstream of the base at position 2, 542, which is a first base of the initiation codon of the downstream gene among the base sequences represented in SEQ ID NO: 19, and the other having a base sequence downstream of the base at position 3, 543, which is a third base of the termination codon of the downstream gene, such that the primer strands extend in opposite directions to each other. Those three combinations provide DNA fragments each containing one of the entire gene group, upstream gene, and downstream gene. Furthermore, there is also a useful combination of primers, which are prepared such that base sequences having over ten or several tens of bases are provided on both ends of the base sequence represented by SEQ ID NO: 17, which extend in opposite directions to each other. This combination allows the production of DNA that corresponds to the base sequence represented by SEQ ID NO: 17, so that a gene corresponding to the downstream gene of the present invention can be produced. Similarly, there is also a useful combination of primers, which are prepared such that base sequences having over ten or several tens of bases are provided on both ends of the base sequence represented by SEQ ID NO: 18, extending in opposite directions to each other. This combination allows the production of DNA that corresponds to the base sequence represented by SEQ ID NO: 18, so that a gene corresponding to the upstream gene of the present invention can be produced.

[0038] A procedure for obtaining genes using those primers is not specifically limited. However, the polymerase chain reaction (PCR) can be most convenient. The reaction conditions are not specifically limited as far as the DNA synthetic reaction produces a reaction product. Conventionally, the reaction may be performed by combining appropriate conditions of a denature temperature of generally 90°C to 100°C, preferably 94°C to 98°C, an annealing temperature of 30°C to 70°C, preferably 37°C to 65°C, more preferably 5°C higher than  $T_m$  of the primer, and an extension temperature of 65°C to 75°C, preferably 72°C. The number of reaction cycles may be usually selected from about 15 to 50 cycles even though the reaction can be repeated until the desired amount of the product will be obtained. The sequence of the gene obtained may be one of the closely-related variants having their own portions different from each other as a

result of the sequence of the DNA strand used as a template and the strength of proof-reading function of DNA polymerase used in the synthesis (the mechanism by which a base incorporated by mistake at the time of DNA replication is removed by the 5' to 3' exonuclease activity of DNA polymerase). However, the closely-related reductase genes can be used in the present invention just as in the case of the original reductase gene used as an origin for primer designing.

[0039] These genes are introduced into the host organisms such that the genes can be expressed in the bodies of the host organisms using expression vectors generally known in the art, allowing the production of organisms each having a high reduction activity enough to produce a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound from an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond by reducing the carbon-carbon double bond. At this time, the downstream gene can obtain a reduction activity when the downstream gene is not used by itself but is combined with the upstream gene.

[0040] Examples of microorganisms for expressing the reductase gene of the present invention are not particularly limited and examples thereof include microorganisms in which host vectors are developed such as bacteria including Escherichia, Bacillus, Pseudomonas, Serratia, Brevibacterium, Corynebacterium, Streptococcus, and Lactobacillus; yeasts such as Saccharomyces, Kluyveromyces, Schizosaccharomyces, Zygosaccharomyces, Yarrowia, Trichosporon, Rhodosporidium, Hansenula, Pichia, and Candida; and fungi such as Neurospora, Aspergillus, Cephalosporium, and Trichoderma. One example of the preferable microorganisms is Escherichia coli. The active microorganism produced does not require any culture medium that contains the above enzyme-inducing substrate as a carbon source. The active microbial cells can be obtained by culturing the cells in a general nutrient culture medium such as an LB medium.

[0041] The reduction reaction using the reduction-active microorganism produced can be performed under the conditions just as in the case of the reaction of the microorganism, from which the present enzyme is derived, disclosed in JP 2000-106891 A.

[0042] In other words, as far as the reducing power of the microorganism can be stably expressed, the reaction for reducing an  $\alpha,\beta$  carbon-carbon double bond of an  $\alpha$ -substituted carbonyl compound having the carbon-carbon double bond may be performed in a culture medium of the microorganism, or performed using cells obtained by the above process, the product obtained by processing the microorganism such as a cell-free extract obtained by disrupting microbial cells cultured by the above process, or the like.

[0043] More specifically, in the case of using the cultured microbial cells, an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond to act as a substrate is added continuously or batchwise to a culture medium in a concentration of 0.1 to 10% by mass, preferably 0.2 to 2% by mass, and is then incubated at a culture temperature of 15 to 40°C, preferably 25 to 37°C, thereby producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound in the culture medium.

[0044] Alternatively, the culture obtained by the above method is subjected to centrifugation or the like to collect microbial cells, and the cells are then suspended in an appropriate solution, for example, an aqueous solution such as a diluted pH buffer. Then, the suspension is added with an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond as a substrate continuously or batchwise in a concentration of, for example, 0.1 to 10% by mass, preferably 0.2 to 2% by mass at a reaction temperature of 15 to 50°C, preferably from 25 to 37°C, more preferably 28 to 35°C, while adjusting the reaction pH to 6.0 to 9.0, preferably from 6.5 to 7.3, thereby producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound in the microbial cell suspension. The pH is preferably maintained constant by means of an aqueous buffer such as one containing potassium phosphate or tris/HCl in a concentration of 10 mM to 1 M.

[0045] The timing and rate or frequency of the addition of the  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond may be freely selected as far as the reaction can be completed within the target time.

[0046] In the case of using a processed microbial product, for example, the culture obtained by the above culture method is subjected to centrifugation to collect microbial cells, and then the cells are disrupted by French pressing or the like to obtain a cell-free extract. Then, the cell-free extract is added to a reaction solution containing an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond as a substrate in a concentration of 0.1 to 10% by mass, preferably from 0.2 to 2% by mass, and also containing 10 mM to 1 M of an ingredient effective in maintaining the pH of the reaction solution. Subsequently, a reaction is carried out at a temperature of 15 to 50°C, preferably 28 to 35°C, thereby producing a corresponding  $\alpha$ -substituted- $\alpha,\beta$ -saturated carbonyl compound.

[0047] In the present invention, the reaction may be performed while a substance (e.g., a compound capable of being oxidized by the microorganism used, such as saccharide or organic acid, preferably glucose or L-lactic acid), which is effective in maintaining the activity of reducing an  $\alpha$ -substituted carbonyl compound having an  $\alpha,\beta$ -carbon-carbon double bond by itself or a mixed solution with an  $\alpha$ -chloroacrylic acid is added continuously or batchwise such that the concentration of the substance reaches 0.1 to 10% by mass, preferably 0.2 to 1% by mass during the reaction. The ratio of the  $\alpha$ -chloroacrylic acid to the added substance to be oxidized may be freely selected between 1:1 and 20:1 on a molar basis. The addition of saccharide or organic acid prolongs a reaction time, allows an increase in the concentration of the target product, an  $\alpha$ -halo- $\alpha,\beta$ -saturated carbonyl compound, in the reaction solution and is advanta-

geous for collecting the product by isolation. In particular, for improving a system for efficiently reproducing the reduction type of co-enzyme NADPH using the substance to be oxidized, an appropriate oxidation-reduction enzyme gene, such as a malate dehydrogenase gene or a glutamate dehydrogenase gene, may be introduced into a microorganism so as to be expressed together with the reductase gene to significantly improve productivity. Such a method is disclosed in publications such as JP 61-128895 A and Biotechnol. Genet. Eng. Rev. 6, 221-270 (1988).

[0048] The reaction may be carried out either in an aerobic or anaerobic environment when the bacterial cells are not in culture. The ratio of the bacterial cells or cell-free extract to the  $\alpha$ -substituted halocarbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond as the substrate, or the timing and rate or frequency of addition of the substrate may be freely selected as far as the reaction can be completed within the desired time.

[0049] In the present invention, an  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -saturated carbonyl compound produced by the reduction of an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond is a metabolic intermediate for the microorganism used and may be further decomposed. In such a case, the decomposition reaction may be terminated by selecting or preparing a host microorganism having no decomposition activity.

[0050] Furthermore, cells or cell-free extract of the microorganism for use in the present invention may be used by fixing the cells or extract to an immobilizing support of various types by a commonly known method such as adsorption, inclusion, or cross-linking. The supports to be used include, but not specifically limited to, polysaccharide-based materials such as cellulose, polymer-based materials, and protein-based materials such as collagen.

[0051] The  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -saturated carbonyl compound produced according to the present invention may be isolated and purified using an ordinary purification method such as solvent extraction or distillation. For example,  $\alpha$ -chloropropionic acid produced from  $\alpha$ -chloroacrylic acid may be obtained by subjecting the culture or reaction solution to organic solvent extraction, distillation, or the like. Furthermore, although an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond is a molecule prochiral at the  $\alpha$ -position, the purity of an enantiomer of the  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -saturated carbonyl compound produced by the reducing method of the present invention, which is a chiral compound, can be determined by means of GC or HPLC with a chiral column or by means of a polarimeter.

[0052] As described above, the present invention provides a group of reductases useful for producing a corresponding  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -saturated carbonyl compound having an S absolute configuration from an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond by reducing the carbon-carbon double bond and also provides a group of genes of the reductases. Furthermore, the present invention provides a manufacturing process using a high-productive organism obtained by the use of those genes.

### BEST MODE FOR CARRYING OUT THE INVENTION

[0053] Hereinafter, the present invention will be described in greater detail with reference to the examples. However, the present invention should not be construed as being limited to these examples. In the examples, all of the base sequence determination does not incorporate a PCR product in a plasmid and the PCR product is directly used as a template. Under standard reaction, isolation, and analysis conditions of the DNA sequencer Model 377 (manufactured by ABI Co., Ltd.), both sequences are decoded.

#### EXAMPLE 1: Detection of Activity of Reducing $\alpha$ -Halo-Carbonyl Compound Having $\alpha$ , $\beta$ -Carbon-Carbon Double Bond

[0054] The reduction activity of the compound was detected using  $\alpha$ -chloroacrylic acid or  $\alpha$ -chloro- $\alpha$ ,  $\beta$ -butenoic acid as a substrate by quantitative determination of a reduction product thereof,  $\alpha$ -chloropropionic acid or  $\alpha$ -chlorobutylic acid with gas chromatography. In addition, 0.4 ml of a reaction solution from which microbial cells were removed by centrifugation or the supernatant of a culture medium was mixed with 0.4 ml of 2N HCl and the resulting mixture was then subjected to a gas chromatographic analysis under the following conditions.

Apparatus: GC-7A (manufactured by Shimadzu Corporation)

Column: Thermon-3000/SHINCARBON A, 2.6 mm x 2.1 m

Carrier gas: nitrogen, 50 ml/min

Detection: FID

Column temperature: 200°C (constant)

Injection: 2 to 10  $\mu$ l, 260°C

Recording: CHROMATOCODER 12 (SIC)

## EXAMPLE 2

(1) Cultivation of Pseudomonas sp. SD811 Strain Using Reduction Substrate as Carbon Source

5 [0055] Pseudomonas sp. SD811 strain was incubated in a culture medium containing the following ingredients:  $\alpha$ -chloroacrylic acid (2 g/l), yeast extract (0.5 g/l), ammonium sulfate (2 g/l), sodium dihydrogen phosphate (1 g/l), dipotassium hydrogen phosphate (1 g/l), and magnesium sulfate (0.1 g/l).

[0056] The medium was prepared as follows.

10 [0057] All of the ingredients mentioned above, except  $\alpha$ -chloroacrylic acid and magnesium sulfate, were dissolved in 950 ml of water. The solution obtained was adjusted to a pH of 7.0, and was then poured into a 5-liter flask and sterilized at 121°C for 20 minutes. Subsequently, after the temperature of the medium decreased to about 70°C, a solution prepared by dissolving  $\alpha$ -chloroacrylic acid and magnesium sulfate in 50 ml of water was adjusted to a pH of 7.0, sterilized through a sterilization filter, and mixed with the medium prepared above. Without oxygen supply or pH adjustment any more, a 5% seed culture (OD 660 nm = 1.10) was inoculated in the medium and the microbial strain

15 was incubated at 30°C for 12 to 24 hours while being shaken.

(2) Cultivation of Pseudomonas sp. SD811 Strain Using Reduction Product as Carbon Source

20 [0058] Pseudomonas sp. SD811 strain was incubated in a culture medium containing the following ingredients: L-lactic acid (2 g/l), yeast extract (0.5 g/l), ammonium sulfate (2 g/l), sodium dihydrogen phosphate (1 g/l), dipotassium hydrogen phosphate (1 g/l), and magnesium sulfate (0.1 g/l).

[0059] The medium was prepared as follows.

25 [0060] All of the ingredients mentioned above, except L-lactic acid and magnesium sulfate, were dissolved in 950 ml of water. The solution obtained was adjusted to a pH of 7.0, and was then poured into a 5-liter flask and sterilized at 121°C for 20 minutes. Subsequently, after the temperature of the medium decreased to about 70°C, a solution prepared by dissolving L-lactic acid and magnesium sulfate in 50 ml of water was adjusted to a pH of 7.0, sterilized through a sterilization filter, and mixed with the medium prepared above. Without oxygen supply or pH adjustment any more, a 5% seed culture (OD 660 nm = 1.10) was inoculated in the medium and the microbial strain was incubated at

30 30°C for 12 to 24 hours while being shaken.

EXAMPLE 3: Cell Suspension Reaction Using  $\alpha$ -Chloroacrylic Acid as Substrate

[0061] In Example 2, two cultures of Pseudomonas sp. SD811 strain cultivated using two different carbon sources were independently centrifuged to collect the microbial cells. Then, the microbial cells were suspended in 20 ml of a solution (adjusted to a pH of 7.3) containing 0.2% of  $\alpha$ -chloroacrylic acid and 100 mM of phosphate buffer (pH 7.3), and the suspension was then reacted at 28°C while being shaken.

35

[0062] From the reaction solution, 0.5 ml was sampled at a specific time and the sample was centrifuged to remove the microbial cells. After that, 0.4 ml of the supernatant from which the microbial cells were removed by centrifugation and 0.1 ml of 6NHC1 were mixed together and then the product was extracted with 0.4 ml of ethyl acetate. The sample

40 extracted was analyzed by the method described in Example 1.

[0063] As a result, in the reaction of microbial cells incubated using a reduction substrate, in association with the consumption of  $\alpha$ -chloroacrylic acid immediately after the reaction, a peak appeared at the position of  $\alpha$ -chloropropionic acid. The reaction rate varied in a nearly linear fashion until the entire substrate was consumed. On the other hand, in the case of the reaction of microbial cells incubated with lactic acid as a carbon source, neither the consumption of  $\alpha$ -chloroacrylic acid nor the peak of  $\alpha$ -chloropropionic acid was detected immediately after the initiation of the reaction.

45 However, after about 5 hours from the initiation, the reduction activity of the microbial cells gradually increased. Fig. 1 shows the results.

## EXAMPLE 4

(1) Cultivation of Burkholderia sp. SD816 Strain Using Reduction Substrate as Carbon Source

50 [0064] Burkholderia sp. SD816 strain was incubated in a culture medium containing the following ingredients:  $\alpha$ -chloroacrylic acid (2 g/l), yeast extract (0.5 g/l), ammonium sulfate (2 g/l), sodium dihydrogen phosphate (1 g/l), dipotassium hydrogen phosphate (1 g/l), and magnesium sulfate (0.1 g/l).

[0065] The medium was prepared as follows.

[0066] All of the ingredients mentioned above, except  $\alpha$ -chloroacrylic acid and magnesium sulfate, were dissolved in 950 ml of water. The solution obtained was adjusted to a pH of 7.0, and was then poured into a 5-liter flask and

sterilized at 121°C for 20 minutes. Subsequently, after the temperature of the medium decreased to about 70°C, a solution prepared by dissolving  $\alpha$ -chloroacrylic acid and magnesium sulfate in 50 ml of water was adjusted to a pH of 7.0, sterilized through a sterilization filter, and mixed with the medium prepared above.

[0067] Without oxygen supply or pH adjustment any more, a 5% seed culture (OD 660 nm = 1.10) was inoculated in the medium and the microbial strain was incubated at 30°C for 12 to 24 hours while being shaken.

## (2) Cultivation of *Burkholderia* sp. SD816 Strain Using Sugar as Carbon Source

[0068] *Burkholderia* sp. SD816 strain was incubated in a culture medium containing the following ingredients: D-glucose (2 g/l), yeast extract (0.5 g/l), ammonium sulfate (2 g/l), sodium dihydrogen phosphate (1 g/l), dipotassium hydrogen phosphate (1 g/l), and magnesium sulfate (0.1 g/l).

[0069] The medium was prepared as follows.

[0070] All of the ingredients mentioned above, except D-glucose and magnesium sulfate, were dissolved in 950 ml of water. The solution obtained was adjusted to a pH of 7.0, and was then poured into a 5-liter flask and sterilized at 121°C for 20 minutes. Subsequently, after the temperature of the medium decreased to about 70°C, a solution prepared by dissolving D-glucose and magnesium sulfate in 50 ml of water was adjusted to a pH of 7.0, sterilized through a sterilization filter, and mixed with the medium prepared above.

[0071] Without oxygen supply or pH adjustment any more, a 5% seed culture (OD 660 nm = 1.10) was inoculated in the medium and the microbial strain was incubated at 30°C for 12 to 24 hours while being shaken.

## EXAMPLE 5: Cell Suspension Reaction Using $\alpha$ -Chloro- $\alpha,\beta$ -Butenoic Acid as Substrate

[0072] In Example 4, two cultures of *Burkholderia* sp. SD816 strain cultivated using two different carbon sources were independently centrifuged to collect the microbial cells. Then, the microbial cells were suspended in 20 ml of a solution (adjusted to a pH of 7.3) containing 0.2% of  $\alpha$ -chloro- $\alpha,\beta$ -butenoic acid and 100 mM of phosphate buffer (pH 7.3), and the suspension was then reacted at 28°C while being shaken.

[0073] From the reaction solution, 0.5 ml was sampled at a specific time and the sample was centrifuged to remove the microbial cells. After that, 0.4 ml of the supernatant from which the microbial cells were removed by centrifugation and 0.1 ml of 6N HCl were mixed together and then the product was extracted with 0.4 ml of ethyl acetate. The sample extracted was analyzed by the method described in Example 1.

[0074] As a result, in the reaction of microbial cells cultivated using a reduction substrate, in association with the consumption of  $\alpha$ -chloro- $\alpha,\beta$ -butenoic acid immediately after the reaction, a peak appeared at the position of  $\alpha$ -chlorobutyric acid. The reaction rate varied in a nearly linear fashion until the entire substrate was consumed. On the other hand, in the case of the reaction of microbial cells incubated with lactic acid as a carbon source, neither the consumption of  $\alpha$ -chloro- $\alpha,\beta$ -butenoic acid nor the peak of  $\alpha$ -chlorobutyric acid was detected immediately after the initiation of the reaction. However, after about 6 to 10 hours from the initiation, the reduction activity of the microbial cells gradually increased. Fig. 2 shows the results.

## EXAMPLE 6: Analysis of Protein Production Pattern with Two-Dimensional Electrophoresis

### (1) Preparation of Crude Enzyme Extract

[0075] The cultures of *Pseudomonas* sp. SD811 strain, having different reduction activities confirmed in Example 2, were incubated for 18 hours. Then, the microbial cells were collected from each culture by centrifugation. The microbial cells collected were washed with sterilized water, followed by resuspending in 50 mM phosphate buffer (pH 7.5). The microbial cells were broken by a BIOMC 7500 ULTRASONIC PROCESSOR (pulsed, 50 of % duty cycle, about 4.5 of output control) and then unbroken cells and insoluble matters were removed by centrifugation (16,400 x g, 5 min, 4°C).

[0076] Similarly, prepared were crude enzyme extracts of the *Burkholderia* sp. SD816 strain cultures having different reduction activities confirmed in Example 5.

### (2) Primary Electrophoresis: Isoelectric Focusing

[0077] A mixture solution was prepared by mixing 1.92 g of urea, 0.53 ml of a 30% acrylamide mixture solution (29.2% (w/v) acrylamide, 0.8% (w/v) N-N'-methylene-bisacrylamide), and 1.0 ml of deionized water. After the urea was completely dissolved in the solution, 0.8 ml of 10% Nonidet P-40, 200  $\mu$ l of Biolight 3/10 Ampholight (BIO-RAD), 8  $\mu$ l of 10% ammonium persulfate, and 5.6  $\mu$ l of TEMED were mixed in the solution. Subsequently, the resulting mixture solution was quickly poured into a glass tube (13 mm in length and 2 mm in inner diameter) having a sealed end, and then a 8 M urea solution was layered on the solution, followed by leaving the mixture untouched for 1 to 2 hours to

make a solidified gel.

[0078] The gel prepared was placed on a semi-micro dry gel electrophoresis apparatus (KS-8110, manufactured by ORIENTAL INSTRUMENTS LTD.), and then a 20 mM sodium hydroxide solution and a 10 mM sulfuric acid solution were poured in upper and lower electrophoresis layers, respectively. Subsequently, the apparatus was pre-activated at 200 V for 15 minutes, 300 V for 15 minutes, and 400 V for 30 minutes.

[0079] The sodium hydroxide solution was removed from the upper electrophoresis layer and the upper side of the gel and then a sample solution (prepared by mixing 100 to 300 µg/12.5 µl of protein in solution, 3 µl of 10% Nonidet P-40, 1.5 µl of Biolight 3/10 Ampholight (BIO-RAD), and 1.5 µl of 2-mercaptoethanol) was placed on the upper side of the gel through a syringe. Subsequently, 20 µl of a sample overlay solution (prepared by mixing 0.48 g of urea, 200 µl of 10% Nonidet P-40, 50 µl of Biolight 3/10 Ampholight (BIO-RAD), and 380 ml of deionized water), and a 20 mM sodium hydroxide solution (appropriate amount) were layered on the gel. Then, the upper electrophoresis layer was filled with a 20 mM sodium hydroxide solution, followed by electrophoresing at 400 V for 12 hours and then at 800 V for 1 hour.

[0080] After the completion of the primary electrophoresis, the gel was removed from the glass tube and then subjected to shaking in 40 ml of deionized water for 5 minutes at room temperature, followed by shaking in 4 ml of an equilibrating buffer (0.5 ml of 0.5 M Tris-HCl (pH 6.8), 1.6 ml of 10% SDS, 0.05 ml of 0.1% BPB, 2 ml of 2-mercaptoethanol, and 1.65 ml of deionized water were mixed) for 20 minutes at room temperature.

### (3) Secondary Electrophoresis: SDS-PAGE

[0081] Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed by a conventional method using a slab gel device (KS-8000 SE type, MARYSOL). More specifically, an equilibrated gel was fixed on the upper end of a 12.5% SDS-PAGE slab gel using 0.5% agarose and then electrophoresis was carried out at a constant current of 25 mA for about 4 hours.

### (4) Detection of Protein: CBB Staining

[0082] The detection of proteins in the slab gel after the electrophoresis was performed by ordinary CBB staining. More specifically, the gel was stained for 1 hour in a CBB solution (prepared by dissolving 0.25 g of Coomassie brilliant blue R-250 in 500 ml of methanol, 50 ml of acetic acid, and 450 ml of deionized water), followed by washing with deionized water and then decolorizing for day and night in a decolorizing solution (50 ml of methanol, 70 ml of acetic acid, and 880 ml of deionized water).

[0083] After that, the gel was immersed in a storage solution (23 ml of 87% (w/v) glycerol solution, 150 ml of ethanol, and 327 ml of deionized water) for 3 hours.

[0084] Comparing the two-dimensional electrophoresis separation patterns of crude enzyme extract samples prepared from the microbial cells in L-lactate culture and microbial cells in  $\alpha$ -chloroacrylic culture of *Pseudomonas* sp. SD811 strain, and the microbial cells in D-glucose culture and microbial cells in  $\alpha$ -chloroacrylic acid culture of *Burkholderia* sp. SD816 strain, a number of product proteins specific to the bacterial cells in  $\alpha$ -chloroacrylic acid culture were found on the nearly same positions in each set of the respective strains. Fig. 3 shows the results of *Pseudomonas* sp. SD811 strain.

### EXAMPLE 7(1): Determination of Terminal Sequence

[0085] For analyzing the proteins specific to the microbial cells in  $\alpha$ -chloroacrylic acid culture, which were found in Example 6, proteins isolated by secondary electrophoresis from the sample of microbial cells in  $\alpha$ -chloroacrylic acid culture of *Burkholderia* sp. SD816 strain were transferred on a PVDF membrane (Immobilon TM Transfermembranes, pore size: 0.45µm, MILLIPORE) using a semi-dry transfer device (TRANS-BLOT R SD Semi-dry Electrophoretic Transfer Cell (Bio-Rad)).

[0086] The transfer was performed according to the standard instructions of the device at a limiting current of 0.8 mA, 13 V, 0.22 to 0.26A for 45 minutes. After the completion of the transfer, the PVDF membrane was stained with the CBB solution. Subsequently, spots corresponding to three different kinds of proteins specifically appeared on the sample of microbial cells in  $\alpha$ -chloroacrylic acid culture were cut out and analyzed on a peptide sequencer (Model 491 Procise (Applied Biosystems)). The result showed that one kind of the proteins was a well-known enzyme, dehalogenase (L-DEX), while the remaining two kinds of the proteins had novel peptide sequences represented by SEQ ID. NOS: 1 and 3.

## EXAMPLE 7(2): Determination of Internal Sequence

[0087] For acquiring further sequence information, two kinds of novel proteins shown in Example 7 were subjected to in-gel partial digestion using lysylendopeptidase.

[0088] After the two-dimensional electrophoresis in Example 6, portions corresponding to two target spots were cut out of the CBB-stained gel. Then, a Tris-buffer containing lysylendopeptidase was added to such a gel section to digest the gel section overnight at 35°C. After that, the reaction solution was subjected to reversed-phase HPLC under the following conditions to isolate fragmented peptides.

Column: TSK gel ODS-120T  
Solvent: TFA/Acetonitrile system, gradient elution  
Flow rate: 1.0 ml/min  
Detection wavelength: 210 nm

[0089] From the resulting chromatogram, an appropriate peak was selected and the fraction thereof was analyzed using a peptide sequencer (Model 491 Procise (Applied Biosystems)). As a result, three internal amino acid sequences represented by SEQ ID NOS: 2, 4, and 5 were obtained.

## EXAMPLE 8: Obtaining N-terminal Portion Gene Fragment of CAA43

[0090] At first, degenerate primers 1 and 2 were designed on the basis of the N-terminal amino acid sequence of CAA43 and the internal amino acid sequence described in SEQ ID NOS: 1 and 2, respectively.

[0091] For the extraction of chromosomal DNA from *Burkholderia* sp. SD816 strain, QIAGEN genomic-tip 100/G and QIAGEN Genomic DNA buffer set (each manufactured by QIAGEN) were used.

[0092] Using BIO-RAD iCycler (manufactured by BIO-RAD), PCR was carried out under the following conditions.

[Composition of Reaction Solution]

[0093]

Chromosomal DNA of <i>Burkholderia</i> sp. SD816	5 ng
Primer 1 (corresponding to SEQ ID NO: 1)	10 pmol
Primer 2 (corresponding to SEQ ID NO: 2)	10 pmol
TaKaRa LATaq	0.5 unit
dNTP mixture (2.5 mM each)	2.0 µl
10 x LA PCR Buffer II (Mg <sup>2+</sup> free)	2.5 µl
25 mM MgCl <sub>2</sub>	2.5 µl
Sterilized distilled water	adjusted to 25 µl

[Reaction Cycle]

[0094]

1 cycle:

Denaturation (95°C, 4 min),  
Annealing (47.9°C, 1 min), and  
Extension (72°C, 2 min).

2 to 30 cycles:

Denaturation (95°C, 1 min),  
Annealing (47.9°C, 1 min), and  
Extension (72°C, 2 min).

[0095] A DNA fragment (350 bp), which might encode a part of the CAA43 gene, was obtained by PCR using the chromosomal DNA of *Burkholderia* sp. SD816 strain as a template. The sequence of the partial fragment was repre-

sented by SEQ ID NO: 11.

#### Example 9: Obtaining Gene Fragment Encoding C-terminal Region of CAA43

5 [0096] Two downstream primers described in SEQ ID NOS: 8 and 9 were designed according to the base sequence represented by SEQ ID NO: 11 obtained in Example 8. The cloning of a gene encoding the C-terminal side of CAA43 was tried using those primers and a TaKaRa LA PCR in vitro Cloning Kit. A reaction or the like was conducted according to the standard instructions attached to the kit. As a result, a DNA fragment (1.3 kb) was obtained by PCR using the chromosomal DNA of *Burkholderia* sp. SD816 strain treated with XbaI as a template, and was then sequenced. The  
10 resulting base sequence was represented by SEQ ID NO: 12 and also a stop codon was identified in this sequence.

#### EXAMPLE 10: Obtaining Genes on N-Terminal Region of CAA43 and Upstream Thereof

15 [0097] A primer for inverted PCR described in SEQ ID NO: 10 was designed according to the base sequence represented by SEQ ID NO: 11 obtained in Example 8 (see "Basics for Genome Engineering", TOKYO KAGAKU DOJIN CO., LTD. (2002)). This primer was combined with the primer described in SEQ ID NO: 8. Then, the inverted PCR was carried out using the chromosomal DNA of *Burkholderia* sp. SD816 strain treated with SalI as a template under the following conditions.

20 [Composition of Reaction Solution]

[0098]

25

30

SalI treated product of SD816 strain chromosomal DNA	200 ng
Primer 1 (SEQ ID NO: 8)	10 pmol
Primer 2 (SEQ ID NO: 10)	10 pmol
TaKaRa LATaq	2.5 units
dNTP mixture (2.5 mM each)	8.0 $\mu$ l
10 x LA PCR Buffer II (Mg <sup>2+</sup> free)	5.0 $\mu$ l
25 mM MgCl <sub>2</sub>	5.0 $\mu$ l
Sterilized distilled water	adjusted to 50 $\mu$ l

[Reaction Cycle]

35

[0099]

1 cycle:

40

Denaturation (94°C, 4.5 min),  
Annealing (55°C, 30 sec), and  
Extension (72°C, 4 min).

2 to 30 cycles:

45

Denaturation (94°C, 30 sec),  
Annealing (55°C, 30 sec), and  
Extension (72°C, 4 min).

50

[0100] As a result, a DNA fragment of about 1.3 kb was obtained. The base sequence of this fragment was represented by SEQ ID NO: 13. The fragment includes the 0.5 kb amino acid sequence of the N-terminal region of CAA43 and a portion encoding the sequence of CAA67 represented by SEQ ID NO: 4. The inventors found that the fragment includes a 0.8 kb portion which may encode the amino acid sequence of the C-terminal region of CAA67. The coding region of CAA67 resides sequentially on the upstream of the coding region of CAA43. Therefore, the inventors found  
55 that both genes form clusters.



## EP 1 473 368 A1

### EXAMPLE 11: Obtaining DNA Fragment Encoding N-Terminal of CAA67

[0101] Two primers on the upstream of CAA67 gene described in SEQ ID NOS: 14 and 15 were designed according to the base sequence revealed in Example 10 encoding the internal amino acid sequence of CAA67. The cloning of a gene encoding the N-terminal side CAA67 was tried using those primers and the TaKaRa LA PCR in vitro Cloning Kit. A reaction or the like was conducted according to the standard instructions attached to the kit. As a result, a DNA fragment (1.8 kb) was obtained by PCR using the chromosomal DNA of *Burkholderia* sp. SD816 strain treated with PstI as a template, and was then sequenced. The resulting base sequence was represented by SEQ ID NO: 12. The inventors confirmed that the DNA fragment was one encoding the internal amino acid sequence of CAA67 described in SEQ ID NOS: 4 and 5 and the DNA fragment encoding the N-terminal amino acid sequence of CAA67 described in SEQ ID NO: 3.

### EXAMPLE 12: Determination of Entire Sequence of Each Gene and Gene Cluster Sequence

[0102] From DNA fragments obtained in Examples 8, 9, and 10, the entire base sequence of CAA43 gene described in SEQ ID NO: 17 was determined using automatic connection-of-nucleic-acid-sequences software (GENETYX-WIN/ATSQ). Similarly, the entire base sequence of CAA67 gene described in SEQ ID NO: 18 was determined using the DNA fragments obtained in Examples 10 and 11. Furthermore, a cluster base sequence described in SEQ ID NO: 19 containing both genes was determined using the DNA fragments obtained in Examples 8 to 11. SEQ ID NOS: 20 and 21 are amino acid sequences corresponding to SEQ ID NOS: 17 and 18, respectively.

### EXAMPLE 13 : Preparation of Gene Fragment Containing CAA43 and CAA67

[0103] Primers described in SEQ ID NOS: 22 and 23 were designed according to the base sequence represented by SEQ ID NO: 19 obtained in Example 12. Then, those primers were combined together and subjected to PCR using the chromosomal DNA of *Burkholderia* sp. SD816 strain as a template under the following conditions to prepare a 2,913 bp DNA fragment encoding the whole length of the reductase gene.

[Composition of Reaction Solution] (50  $\mu$ l in total)

[0104]

SD816 strain chromosomal DNA (5 $\mu$ g/ $\mu$ l)	4.0 $\mu$ l
10 $\mu$ M primer 1 (SEQ ID NO: 22)	1.5 $\mu$ l
10 $\mu$ M primer 2 (SEQ ID NO: 23)	1.5 $\mu$ l
TOYOBO KOD-Plus-(1 unit/ $\mu$ l)	1.0 $\mu$ l
dNTP mixture (2.5 mM each)	5.0 $\mu$ l
10 x KOD PCR Buffer (Mg <sup>2+</sup> free)	5.0 $\mu$ l
25 mM MgCl <sub>2</sub>	2.0 $\mu$ l
Sterilized distilled water	30 $\mu$ l

[Reaction Cycle]

[0105]

1 cycle:

Denaturation (94°C, 2 min),

2 to 30 cycles:

Denaturation (94°C, 15 sec),

Annealing (52.3°C, 30 sec), and

Extension (68°C, 3 min).

## EXAMPLE 14: Construction of CAA43 and CAA67 Expression Systems

[0106] The DNA fragment obtained in Example 13 was inserted into the downstream of T7 promoter in the expression vector pET101/D-TOPO, followed by introducing into *Escherichia coli* BL21(DE3). Ligation between the insert and the vector, transformation, and gene expression were performed using a pET101 Directional TOPO-Expression Kit (Invitrogen).

## EXAMPLE 15: Reduction Reaction Using CAA43 and CAA67 Active Microbial Cells

[0107] The microbial cells obtained in Example 14 were incubated in a 5 ml LB culture medium (1% Bacto Tryptone (DIFCO), 0.5% Bacto Yeast Extract (DIFCO), 1% Sodium chloride (Nacalai Tesque), and 100 mg/ml ampicillin) (37°C, 130 rpm, 10 h.). The resulting cells were suspended in 1 ml of 60 mM phosphate buffer (1mM DTT added, pH 7.1). Then, the microbial cells were disrupted by sonication (BRANSON Digital Sonifier) and then centrifuged (15,000 rpm, 4°C, and 10 min). The reduction activity of the supernatant of the cell-disrupted solution was measured according to the method shown in Example 1. At this time, various co-enzymes were added to the reaction solution and the reduction activity thereof was then measured. Consequently, a sufficient reduction activity was observed only when NADPH (reduced nicotinamide adenine dinucleotide phosphate) was added to the reaction solution.

[0108] Next, 1/10 volume of NADPH was added to a reaction solution (3 mM 2-CAA, 0.65 mM NADPH, 60 mM Ammonium acetate buffer (pH 7.1)) and then the decrease of NADPH over time at the time of reacting at 30°C was measured by variations in absorbance at 339 nm in a cell having an optical path length of 0.2 cm. The enzyme level to decrease 1 mmol of NADPH per minute was defined as an enzymatic activity of 1 unit to calculate a specific activity (units/mg). Table 1 shows the 2-CAA reductase activity of the transformant and that of *E.coli* BL21 (DE3). A significant 2-CAA reduction activity was observed in the transformant.

Table 1

2-CAA reductase activities of transformant and host	
Strain	Specific activity (units/mg)
<i>E. coli</i> BL21 (DE3)	0.06
<i>E. coli</i> BL21 (DE3) pET101/D/67&43	0.92

## INDUSTRIAL APPLICABILITY

[0109] The present invention provide a base sequence encoding a related enzyme having a high catalytic activity useful in producing a corresponding  $\alpha$ -substituted- $\alpha$ ,  $\beta$ -saturated carbonyl compound from an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond by reducing the carbon-carbon double bond using an enzyme produced by a microorganism by a process favored with high profitability, good operability, and excellent processing safety. Furthermore, the present invention provides a reductase and a gene product thereof useful in producing a corresponding highly-purified and optically-active  $\alpha$ -substituted- $\alpha$ , $\beta$ -saturated carbonyl compound, which is useful as chiral building blocks of medical and agricultural chemicals and the like with respect to the  $\alpha$  position, from an  $\alpha$ -substituted carbonyl compound having an  $\alpha$ ,  $\beta$ -carbon-carbon double bond prochiral at the  $\alpha$ -position by hydrogenating the carbon-carbon double bond.

Address	Name	Deposit date	Deposit number
Central 6, Higashi 1-chome 1-1, Tsukuba-shi, Ibaraki prefecture, Japan (Postal code number, and 305-8566)	International Patent Organism Depository National Institute of Advanced Industrial Science Technology, an Independent Administrative Institution	1998/4/2	FERM BP-6767
		1998/4/2	FERM BP-6768
		1998/4/2	FERM BP-6769
		1999/6/28	FERM BP-6770

## SEQUENCE LISTING

5 <110> SHOWA DENKO K. K.

<120> REDUCTASE GENE FOR  $\alpha$ -SUBSTITUTED- $\alpha,\beta$ -UNSATURATED CARBONYL COMPOUND

10 <130> EPA-63516

<150> JP2002-030127

15 <151> 2002-02-06

<150> JP2002-281236

20 <151> 2002-09-26

<160> 23

25 <170> PatentIn version 3.1

<210> 1

30 <211> 16

<212> PRT

<213> Burkholderia sp.

35 <400> 1

40 Val Met Ala Ala Val Ile His Lys Lys Gly Gly Pro Asp Asn Phe Val

1 5 10 15

<210> 2

45 <211> 15

<212> PRT

<213> Burkholderia sp.

50 <400> 2

55 Asp Leu Asp Leu Asp Asp Val His Leu Ala Gly Leu Met Leu Lys

1 5 10 15

5           <210> 3  
           <211> 13  
           <212> PRT  
           <213> Burkholderia sp.  
  
 10          <220>  
           <221> MISC\_FEATURE  
           <222> (1).. (1)  
 15          <223> Unknown  
  
           <220>  
 20          <221> MISC\_FEATURE  
           <222> (12).. (12)  
           <223> Unknown  
  
 25          <400> 3  
  
           Xaa Asp Val Leu Val Thr Asp Val Leu Val Val Xaa Gly  
 30           1                   5                   10  
  
           <210> 4  
 35          <211> 25  
           <212> PRT  
           <213> Burkholderia sp.  
  
 40          <400> 4  
  
           Val Phe Val Asp Phe Arg Glu Thr Lys Pro Glu Glu Trp Ala Pro Asp  
 45           1                   5                   10                   15  
  
           Ser Leu Thr Gly Thr Phe Leu Gly Lys  
 50                   20                   25  
  
           <210> 5  
 55          <211> 30  
           <212> PRT

<213> Burkholderia sp.

<400> 5

Glu Leu Leu Ser Gly Leu Asp Ala Asp Tyr Gly Thr Arg Gly Ser Leu

1 5 10 15

Glu Asp Thr Thr Gly Leu Met Met Glu Phe Ser Ser Thr His

20 25 30

<210> 6

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Forward primer for N-terminal fragment of CAA43

<220>

<221> misc\_feature

<222> (6)..(6)

<223> Inosine

<220>

<221> misc\_feature

<222> (9)..(9)

<223> Inosine

<220>

<221> misc\_feature

<222> (12)..(12)

<223> Inosine

<400> 6

atggcngcng tnatcayaa

20

<210> 7

```

5      <211> 20
      <212> DNA
      <213> Artificial

      <220>
10     <223> Reverse primer for N-terminal fragment of CAA43

      <220>
      <221> misc_feature
15     <222> (3)..(3)
      <223> Inosine

      <220>
20     <221> misc_feature
      <222> (6)..(6)
      <223> Inosine

25     <220>
      <221> misc_feature
      <222> (12)..(12)
      <223> Inosine

30     <400> 7
      ccngcnarrt gnacrtcgtc

      <210> 8
40     <211> 20
      <212> DNA
      <213> Artificial

      <220>
45     <223> Forward primer for C-terminal fragment of CAA43

      <400> 8
50     cgccccctcgg tgcctacagc

      <210> 9
55     ccgctcctcgcg cgcctcctcgcg

```

<211> 20  
 <212> DNA  
 5 <213> Artificial  
  
 <220>  
 10 <223> Reverse primer for C-terminal fragment of CAA43  
  
 <400> 9  
 15 ctaccccgcc gaaaaactga 20  
  
 <210> 10  
 <211> 20  
 20 <212> DNA  
 <213> Artificial  
  
 25 <220>  
 <223> Primer for inverted PCR  
  
 <400> 10  
 30 ccgggcgcgc caaccttaac 20  
  
 <210> 11  
 35 <211> 350  
 <212> DNA  
 <213> Burkholderia sp.  
 40  
 <400> 11  
 ggtaattcat aagaagggtg gaccgataa cttcgatgg gaggaagta aggttggtc 60  
 45 gcccgcccg ggtcaagtgc gactgcgcaa tacggccatc ggggtaaact tcctggatac 120  
 ttatcaccgc gcaggcattc ctacccgct ggtcgttgg gagccgcga ttgtggtcg 180  
 50 cttcgaagcc gccgctgtgg ttgaggaagt cgggccggt gtaaccgact tcaccgttg 240  
 55 tgagcgggtc tgcacttgc ttccgccct cggcgctac agccaggagc gcctctacc 300

cgccgaaaaa ctgatcaagg ttccaaagga cctggatctt gatgacgtgc 350

5 <210> 12  
 <211> 1069  
 <212> DNA  
 10 <213> Burkholderia sp.

<400> 12

15 tcaaggttcc aaaggacctg gatcttgatg acgtgcacct cgcgggattg atgctcaagg 60

ggatgacagc acaatatctg ctgcatcaga cgcacaaggt aaagccgggt gactacgtgt 120

20 tgattcacgc ggcggctggc ggcatgggcc acatcatggt tccttgggcg cgccacctcg 180

gcgctaccgt gatcgggacg gtcagcacgg aagaaaaggc tgagactgct cgcaaacctcg 240

25 gtgcccacca taccatcaat tattccactc aggatcttcg tgaggtagtt ogcgaaatca 300

cgggcgggaa ggggtgtgac gtggtctacg attccatcgg taaagacaca ctccagaagt 360

30 cgctcgactg tctgcgcccg cgcggtatgt gtgcggccta cgggcacgca tccggcgtgg 420

35 cagatccgat cagggtcgtc gaggacctcg gtgtacgtgg atcgtgttc attactagac 480

ccgcactctg gcattacatg tcgaaccgca gtgagattga cgaagggtcg aagtcctgt 540

40 tcgatgccgt caaggcgggt gtactccata gcagtgtcg aaagaccttc cctctgcggg 600

aggcagcggc ggcgcacaaa tacatgggtg gtcgtcagac gatcggctcg attgttttgc 660

45 ttccgcaagc gtaggtagcc gtagggcgtc accccggaat ttcgggtga ccgaaaacgt 720

cgccgtgaac gcctatctgc ggatgttggc ggttgggtgg cattattttt cggcagcccg 780

50 tcgttcgctg ccggctcgta cgttgccgtt ttggacagtt cgtcaacggg gcgattgtga 840

55 ttccaaccc gcgaccctcg attgggtcaa cgtgtgtttg actttctgtg aactctgtct 900



EP 1 473 368 A1

aagtacacga gaggtgtgta gatcagacgc gtgccgaagc tactcggatg ttccctgtcg 960

5 aggcaattga aagagtcaat acacatgaac gtattcaata gagagatagt gctttccgtc 1020

cggcactgga cggataaact tttcagcttt accgcaaccc gtaacgccg 1069

10

<210> 13

<211> 1617

15 <212> DNA

<213> Burkholderia sp.

20 <400> 13

gtcgacttcc gcgaaacgaa accggaggaa tgggcgcctg attcactcac aggcaccttc 60

25 ttgggcaagt gtgtcccgaa tttcatgacc accccggtac aggttgccgc gtcacgcac 120

tacacgatcg gcggtctgaa agtcgatgtg gatggccgta ccaatcttcc gaagctctac 180

30 gctgtcggcg agttggccgg tggcgtgcat ggcgccaacc gccacgggtg cacggcgctg 240

gtcgatgcca tgacgtacgg ccggattgct ggaaggcacg cggcgggaag cctcaacggc 300

35 gcggctgcga cgggaggtgc agcgttcta cccgcaggca gcaaagcggg aaaggcaagc 360

40 cggattgagg gcgcaatgag cgatctgcgc cgcgcaaacc agtcgcctct tggctctatt 420

cgtgatgccg tacggcttca acgcgttggg gagctgtttg ctgaactctt ggacgaggtc 480

45 cgctcggtcg gttggaacgg ctacaaggaa atgcaggaaa tcttgccgct cgagcgtgcg 540

atcaagctgt ctgacgctat gcgccaggcc atgttacgcc gcacggagac acgaggagtc 600

50 cactatcggg ccgatttccc gagctccagt gatgcatggt tgaagaagca ggtatttgca 660

55 ttgcgcgatg gggcgttgca tttcaaagac gttccctct aatcaaagtc gcataagcgc 720

gatattgcat caggtatttg tctcgtgtg gtgtcatttg tgcctcggcg cccaagggtg 780  
 5 agagcggaag gtggagctcg ccgccgcggg cagatctggt gtgtgtcgtt gttccgtatc 840  
 ggtagcaaaa acaatctgac ttcgctagcg ggcaggtaag ccgacggcgt atgctcggcg 900  
 10 aaggttcttg aattgagttc gtggtgacca tgcgctcgc gtgtgaatcg ctatttagga 960  
 gactttatca tggtaatggc agcggtaatt cataagaagg gtggaccgga taacttcgta 1020  
 15 tgggaggaag ttaaggttgg ctgcccggc ccgggtcaag tgcgactgcg caatacggcc 1080  
 atcggggtaa acttcctgga tacttatcac cgcgcaggca ttcctcaccg gctggtcgtt 1140  
 ggcgagccgc cgattgtggt cggcttcgaa gccgccgctg tggttgagga agtcgggtccc 1200  
 25 ggtgtaaccg acttcacogt tggtagcgg gtctgcactt gtcttcggcc cctcgggtgcc 1260  
 tacagccagg agcgcctcta cccgccgaa aaactgatca aggttccaaa ggacctggat 1320  
 30 cttgatgacg tgcacctcgc gggattgatg ctcaagggga tgacagcaca atatctgctg 1380  
 catcagacgc acaaggtaaa gccgggtgac tacgtgttga ttcacgcggc ggctggcggc 1440  
 atgggccaca tcatggttcc ttgggcgcgc caccctggcg ctaccgtgat cgggacggtc 1500  
 40 agcacggaag aaaaggctga gactgctcgc aaactcggct gccaccatac catcaattat 1560  
 tccactcagg atttcgctga ggtagttcgc gaaatcacgg gcgggaaggg tgctcgac 1617  
 45  
 <210> 14  
 <211> 20  
 <212> DNA  
 50 <213> Artificial  
 <220>  
 55 <223> Forword primer for N-terminal fragment of CAA67

<400>	14		
catgaaattc	gggacacact		20
<210>	15		
<211>	20		
<212>	DNA		
<213>	Artificial		
<220>			
<223>	Reverse primer for N-terminal fragment of CAA67		
<400>	15		
agaaggtgcc	tgtgagtga		20
<210>	16		
<211>	1640		
<212>	DNA		
<213>	Burkholderia sp.		
<400>	16		
aagcgctactc	cagcgttagc	tggccaatcc	ttgcgtggag
caccgtgacg	tccactggcg		60
gctcaaacgt	cgttgacttg	cctgtctcga	acacttccga
cacgcgtcc	tgcaactgct		120
tcttccagtc	tgtaatctgg	ttcggatgaa	cgtcgcactg
ctgcgccagt	tctgccagcg		180
tcttgtcgcc	cctcggggct	gccagcccta	tctttgcctt
gaataccgtg	ctatgcgacc		240
ggcggcttct	tttgtcatcg	ttgggtcct	cttgccggtc
cttataccag	ttcagttccc		300
ggccgttcga	cttaccgacc	tgctgaatt	tccggccgca
cctctaacc	aagatcattt		360
ctgacgattt	gtagaagtt	ttctgcgttc	aggccacttt
ttcgccgct	ttccgagggg		420
atagtattgc	gacaaatgtg	agcgatccgt	agccaacggg
tattgcgagg	cggtgccgc		480

EP 1 473 368 A1

ttcggcgggt cacacctatc ctgtgtggtc gcacacaagg ttccgcgacgt caataaagat 540

5 gatttggaga catgcacgtg atgttctccc ttgatgtcta gcggtcgttg aggatcattt 600

aatccaatgt ttgacaggag gaggatgttc atgtcggatg ttcttgtaac agacgtgttg 660

10 gtggtggcgc agggctgcgc aggccaaacc gctgcgctta ctgcaagcga gtcgggttgt 720

gacgtcatca tgcttggaga cggccgcgca cagagcaccg ctgtttccac cggttcctt 780

15 acttatgccg cgcacgaagg tttcaatcgt gccagctct acgaagcga gtcacagacc 840

acaggcaagg gcttgtgtga tgtagcgtc ttgaggcgac ttgtcgatga agtccgaaa 900

gaaatggcgg agttgattga gacatataag gttcctgtcg ataacaccga gcgtggagtg 960

25 agggcgcgcc gggcagtggg taagagcgga aaagagcttc tctccgatt ggacgggat 1020

tacgggacgc gtggttccct caggacacg acgggocctaa tgatggagtt ctcgtcgaca 1080

30 cacggaacag cgctctatgc ccagttgcgt aaagccgtga acacggcgcc aaagattcgg 1140

cggtacgcg gaagtgcgt ggttctcgaa cccggtcca ccacggtcgg tgcgttgtc 1200

gatggcgagc cggtagcat cgcggctcgt tcgatcatct tggcgactgg agggattcag 1260

40 ggctctacg aggtcacgga taaccgcat acgtcacgg gtgatggtca tggcatggcg 1320

atggacgctg gcgcggagtt cgtcgacatg gattcatgc agttctaccc gctttcagtc 1380

45 aatgaggagg gcgcaccgac actcttcttc tatccgatt tcccaggcg cgccaagtc 1440

attgacgacg ggggcccgaac cgtcctggtg aagcatctcg gcgagggtc gcaatacctt 1500

tcggagttgc ataattggga tcagctagcc gcggtgttac agacggagat tgcgaaggc 1560

55 aggaaggtat ttgtcgactt ccgcgaacg aaaccggagg aatgggcgcc tgattcactc 1620

acaggcacct tcttgggcaa

1640

5

&lt;210&gt; 17

&lt;211&gt; 1002

&lt;212&gt; DNA

10

&lt;213&gt; Burkholderia sp.

&lt;220&gt;

15

&lt;221&gt; CDS

&lt;222&gt; (1).. (1002)

&lt;223&gt;

20

&lt;400&gt; 17

atg gta atg gca gcg gta att cat aag aag ggt gga ccc gat aac ttc 48

Met Val Met Ala Ala Val Ile His Lys Lys Gly Gly Pro Asp Asn Phe

25

1 5 10 15

gta tgg gag gaa gtt aag gtt ggc tcg ccc ggc ccg ggt caa gtg cga 96

Val Trp Glu Glu Val Lys Val Gly Ser Pro Gly Pro Gly Gln Val Arg

30

20 25 30

ctg cgc aat acg gcc atc ggg gta aac ttc ctg gat act tat cac cgc 144

Leu Arg Asn Thr Ala Ile Gly Val Asn Phe Leu Asp Thr Tyr His Arg

35

35 40 45

gca ggc att cct cac ccg ctg gtc gtt ggc gag ccg ccg att gtg gtc 192

Ala Gly Ile Pro His Pro Leu Val Val Gly Glu Pro Pro Ile Val Val

40

50 55 60

ggc ttc gaa gcc gcc gct gtg gtt gag gaa gtc ggt ccc ggt gta acc 240

Gly Phe Glu Ala Ala Ala Val Val Glu Glu Val Gly Pro Gly Val Thr

45

65 70 75 80

gac ttc acc gtt ggt gag cgg gtc tgc act tgt ctt ccg ccc ctc ggt 288

Asp Phe Thr Val Gly Glu Arg Val Cys Thr Cys Leu Pro Pro Leu Gly

50

85 90 95

EP 1 473 368 A1

	gcc tac agc cag gag cgc ctc tac ccc gcc gaa aaa ctg atc aag gtt	336
	Ala Tyr Ser Gln Glu Arg Leu Tyr Pro Ala Glu Lys Leu Ile Lys Val	
5	100 105 110	
	cca aag gac ctg gat ctt gat gac gtg cac ctc gcg gga ttg atg ctc	384
10	Pro Lys Asp Leu Asp Leu Asp Asp Val His Leu Ala Gly Leu Met Leu	
	115 120 125	
	aag ggg atg aca gca caa tat ctg ctg cat cag acg cac aag gta aag	432
15	Lys Gly Met Thr Ala Gln Tyr Leu Leu His Gln Thr His Lys Val Lys	
	130 135 140	
20	ccg ggt gac tac gtg ttg att cac gcg gcg gct ggc ggc atg ggc cac	480
	Pro Gly Asp Tyr Val Leu Ile His Ala Ala Ala Gly Gly Met Gly His	
25	145 150 155 160	
	atc atg gtt cct tgg gcg cgc cac ctc ggc gct acc gtg atc ggg acg	528
30	Ile Met Val Pro Trp Ala Arg His Leu Gly Ala Thr Val Ile Gly Thr	
	165 170 175	
	gtc agc acg gaa gaa aag gct gag act gct cgc aaa ctc ggc tgc cac	576
35	Val Ser Thr Glu Glu Lys Ala Glu Thr Ala Arg Lys Leu Gly Cys His	
	180 185 190	
40	cat acc atc aat tat tcc act cag gat ttc gct gag gta gtt cgc gaa	624
	His Thr Ile Asn Tyr Ser Thr Gln Asp Phe Ala Glu Val Val Arg Glu	
	195 200 205	
45	atc acg ggc ggg aag ggt gtc gac gtg gtc tac gat tcc atc ggt aaa	672
	Ile Thr Gly Gly Lys Gly Val Asp Val Val Tyr Asp Ser Ile Gly Lys	
50	210 215 220	
	gac aca ctc cag aag tcg ctc gac tgt ctg cgg ccg cgc ggt atg tgt	720
55	Asp Thr Leu Gln Lys Ser Leu Asp Cys Leu Arg Pro Arg Gly Met Cys	
	225 230 235 240	

EP 1 473 368 A1

	gcg gcc tac ggg cac gca tcc ggc gtg gca gat ccg atc agg gtc gtc	768
	Ala Ala Tyr Gly His Ala Ser Gly Val Ala Asp Pro Ile Arg Val Val	
5	245 250 255	
	 gag gac ctc ggt gta cgt gga tgc ctg ttc att act aga ccc gca ctc	816
10	Glu Asp Leu Gly Val Arg Gly Ser Leu Phe Ile Thr Arg Pro Ala Leu	
	260 265 270	
	 tgg cat tac atg tgc aac cgc agt gag att gac gaa ggg tgc aag tgc	864
15	Trp His Tyr Met Ser Asn Arg Ser Glu Ile Asp Glu Gly Ser Lys Cys	
	275 280 285	
	 ctg ttc gat gcc gtc aag gcg ggt gta ctc cat agc agt gtc gca aag	912
20	Leu Phe Asp Ala Val Lys Ala Gly Val Leu His Ser Ser Val Ala Lys	
	290 295 300	
25	 acc ttc cct ctg cgg gag gca gcg gcg gcg cac aaa tac atg ggt ggt	960
	Thr Phe Pro Leu Arg Glu Ala Ala Ala Ala His Lys Tyr Met Gly Gly	
30	305 310 315 320	
	 cgt cag acg atc ggc tgc att gtt ttg ctt ccg caa gcg tag	1002
	Arg Gln Thr Ile Gly Ser Ile Val Leu Leu Pro Gln Ala	
35	325 330	
	 <210> 18	
	<211> 1644	
40	<212> DNA	
	<213> Burkholderia sp.	
45	 <220>	
	<221> CDS	
	<222> (1).. (1644)	
50	<223>	
	 <400> 18	
55	atg tgc gat gtt ctt gta aca gac gtg ttg gtg gtg ggc gag ggc tgc	48
	Met Ser Asp Val Leu Val Thr Asp Val Leu Val Val Gly Glu Gly Cys	

EP 1 473 368 A1

	1	5	10	15	
5	gca ggc caa acc gct gcg ctt act gca agc gag tcg ggt tgt gac gtc				96
	Ala Gly Gln Thr Ala Ala Leu Thr Ala Ser Glu Ser Gly Cys Asp Val				
		20	25	30	
10	atc atg ctt gga gac ggc cgc gca ccg agc acc gct gtt tcc acc ggc				144
	Ile Met Leu Gly Asp Gly Arg Ala Pro Ser Thr Ala Val Ser Thr Gly				
		35	40	45	
15	ttc ctt act tat gcc gcg cac gaa ggt ttc aat cgt gcc cag ctc tac				192
	Phe Leu Thr Tyr Ala Ala His Glu Gly Phe Asn Arg Ala Gln Leu Tyr				
20		50	55	60	
	gaa gcg atg tca cag acc aca ggc aag ggc ttg tgt gat gta gcg ctc				240
25	Glu Ala Met Ser Gln Thr Thr Gly Lys Gly Leu Cys Asp Val Ala Leu				
	65	70	75	80	
	ttg agg cga ctt gtc gat gaa gct ccg aaa gaa atg gcg gag ttg att				288
30	Leu Arg Arg Leu Val Asp Glu Ala Pro Lys Glu Met Ala Glu Leu Ile				
		85	90	95	
35	gag aca tat aag gtt cct gtc gat aac acc gag cgt gga gtg agg gcg				336
	Glu Thr Tyr Lys Val Pro Val Asp Asn Thr Glu Arg Gly Val Arg Ala				
		100	105	110	
40	cgc cgg gca gtg ggt aag agc gga aaa gag ctt ctc tcc gga ttg gac				384
	Arg Arg Ala Val Gly Lys Ser Gly Lys Glu Leu Leu Ser Gly Leu Asp				
		115	120	125	
45	gcg gat tac ggg acg cgt ggt tcc ctc gag gac acg acg ggc cta atg				432
	Ala Asp Tyr Gly Thr Arg Gly Ser Leu Glu Asp Thr Thr Gly Leu Met				
50		130	135	140	
	atg gag ttc tcg tcg aca cac gga aca gcg ctc tat gcc cag ttg cgt				480
55	Met Glu Phe Ser Ser Thr His Gly Thr Ala Leu Tyr Ala Gln Leu Arg				
	145	150	155	160	



	aaa gcc gtg aac acg gcg cca aag att cgg cgc gta cgc gga agt gcg	528
5	Lys Ala Val Asn Thr Ala Pro Lys Ile Arg Arg Val Arg Gly Ser Ala	
	165 170 175	
10	ctg gtt ctc gaa ccc ggt tcc acc acg gtc ggt gcg ctt gtc gat ggc	576
	Leu Val Leu Glu Pro Gly Ser Thr Thr Val Gly Ala Leu Val Asp Gly	
	180 185 190	
15	gag ccg gtg acg atc gcg gct cgt tcg atc atc ttg gcg act gga ggg	624
	Glu Pro Val Thr Ile Ala Ala Arg Ser Ile Ile Leu Ala Thr Gly Gly	
	195 200 205	
20	att cag ggc ctc tac gag gtc acg gat aac ccg oat acg ctc acg ggt	672
	Ile Gln Gly Leu Tyr Glu Val Thr Asp Asn Pro His Thr Leu Thr Gly	
25	210 215 220	
30	gat ggt cat ggc atg gcg atg gac gct ggc gcg gag ttc gtc gac atg	720
	Asp Gly His Gly Met Ala Met Asp Ala Gly Ala Glu Phe Val Asp Met	
	225 230 235 240	
35	gag ttc atg cag ttc tac ccg ctt tca gtc aat gag gag ggc gca ccg	768
	Glu Phe Met Gln Phe Tyr Pro Leu Ser Val Asn Glu Glu Gly Ala Pro	
	245 250 255	
40	aca ctc ttc ttc tat ccg gat ttc ccc agg cgc gcc aag ctc att gac	816
	Thr Leu Phe Phe Tyr Pro Asp Phe Pro Arg Arg Ala Lys Leu Ile Asp	
	260 265 270	
45	gac ggg ggc cga aac gtc ctg gtg aag cat ctc ggc gag ggc tcg caa	864
	Asp Gly Gly Arg Asn Val Leu Val Lys His Leu Gly Glu Gly Ser Gln	
50	275 280 285	
55	tac ctt tog gag ttg cat aat tgg gat cag cta gcc gcg gtg gta cag	912
	Tyr Leu Ser Glu Leu His Asn Trp Asp Gln Leu Ala Ala Val Val Gln	
	290 295 300	

EP 1 473 368 A1

	acg gag att gtc gaa ggc agg aag gta ttt gtc gac ttc cgc gaa acg	960
	Thr Glu Ile Val Glu Gly Arg Lys Val Phe Val Asp Phe Arg Glu Thr	
5	305                      310                      315                      320	
	aaa ccg gag gaa tgg gcg cct gat tca ctc aca ggc acc ttc ttg ggc	1008
10	Lys Pro Glu Glu Trp Ala Pro Asp Ser Leu Thr Gly Thr Phe Leu Gly	
	325                      330                      335	
	aag tgt gtc ccg aat ttc atg acc acc ccg gta cag gtt gcg ccg tca	1056
15	Lys Cys Val Pro Asn Phe Met Thr Thr Pro Val Gln Val Ala Pro Ser	
	340                      345                      350	
	tcg cac tac acg atc ggc ggt ctg aaa gtc gat gtg gat ggc cgt acc	1104
20	Ser His Tyr Thr Ile Gly Gly Leu Lys Val Asp Val Asp Gly Arg Thr	
	355                      360                      365	
	aat ctt ccg aag ctc tac gct gtc ggc gag ttg gcc ggt ggc gtg cat	1152
25	Asn Leu Pro Lys Leu Tyr Ala Val Gly Glu Leu Ala Gly Gly Val His	
	370                      375                      380	
30		
	ggc gcc aac cgc cac ggt ggc acg gcg ctg gtc gat gcc atg acg tac	1200
	Gly Ala Asn Arg His Gly Gly Thr Ala Leu Val Asp Ala Met Thr Tyr	
35	385                      390                      395                      400	
	ggc cgg att gct gga cgg cac gcg gcg gga agc ctc aac ggc gcg gct	1248
40	Gly Arg Ile Ala Gly Arg His Ala Ala Gly Ser Leu Asn Gly Ala Ala	
	405                      410                      415	
	gcg acg gga ggt gca gcg ctt cta ccc gca ggc agc aaa gcg gga aag	1296
45	Ala Thr Gly Gly Ala Ala Leu Leu Pro Ala Gly Ser Lys Ala Gly Lys	
	420                      425                      430	
	gca agc cgg att gag ggc gca atg agc gat ctg cgc cgc gca aac cag	1344
50	Ala Ser Arg Ile Glu Gly Ala Met Ser Asp Leu Arg Arg Ala Asn Gln	
	435                      440                      445	
55		
	ctc gct ctt ggt cct att cgt gat gcc gta cgg ctt caa cgc gtt ggg	1392

EP 1 473 368 A1

Leu Ala Leu Gly Pro Ile Arg Asp Ala Val Arg Leu Gln Arg Val Gly  
450 455 460

5

gag ctg ttt gct gaa ctc ttg gac gag gtc cgc tcg ttc ggt tgg aac 1440  
Glu Leu Phe Ala Glu Leu Leu Asp Glu Val Arg Ser Phe Gly Trp Asn  
10 465 470 475 480

ggc tac aag gaa atg cag gaa atc ttg cgc gtc gag cgt gcg atc aag 1488  
15 Gly Tyr Lys Glu Met Gln Glu Ile Leu Arg Val Glu Arg Ala Ile Lys  
485 490 495

ctg tct gac gct atg cgc cag gcc atg tta cgc cgc acg gag aca cgc 1536  
20 Leu Ser Asp Ala Met Arg Gln Ala Met Leu Arg Arg Thr Glu Thr Arg  
500 505 510

gga gtc cac tat cgg gcc gat ttc ccg agc tcc agt gat gca tgg ttg 1584  
25 Gly Val His Tyr Arg Ala Asp Phe Pro Ser Ser Ser Asp Ala Trp Leu  
515 520 525

aag aag cag gta ttt gca ttg cgc gat ggg gcg ttg cat ttc aaa gac 1632  
30 Lys Lys Gln Val Phe Ala Leu Arg Asp Gly Ala Leu His Phe Lys Asp  
530 535 540

gtt ccc ctc taa 1644  
Val Pro Leu  
40 545

<210> 19  
45 <211> 3938  
<212> DNA  
<213> Burkholderia sp.

50 <400> 19  
aagcgtactc cagcgttagc tgccaatcc ttgcgtggag caccgtgacg tccactggtg 60

55 gctcaaacgt cgttgacttg cctgtctega acacttccga cagcgctcc tgcaactgct 120

tcttccagtc tgtaatctgg ttcggatgaa cgtcgactg ctgcgccagt tctgccagcg 180  
 5 tcttgcgcc cctcggggct gccagcccta tctttgcctt gaataccgtg ctatgcgacc 240  
 ggccgcttct tttgtcatcg ttgggtcctt cttgccggtc cttataccag ttcagttccc 300  
 10 ggccgttcga cttaccgacc tgcctgaatt tccggccgca cctctaacc aagatcattt 360  
 ctgacgattt ggtagaagtt ttctgcgttc agggcacttt ttccggccgt ttcggagggg 420  
 15 atagtattgc gacaaatgtg agcgatccgt agccaacggg tattgcgagg cggctgccgc 480  
 ttccggcgggt cacacctatc ctgtgtgtgc gcacacaagg ttccgcgacgt caataaagat 540  
 20 gatttgaga catgcacgtg atgttctccc ttgatgtcta gcggtcgtt aggatcattt 600  
 25 aatccaatgt ttgacaggag gaggatgttc atgtcggatg ttcttgaac agcgtgttg 660  
 gtggtggcg agggctgcgc agggcaaacc gctgcgtta ctgaagcga gtcgggttgt 720  
 30 gacgtcatca tgcttgaga cggccgcgca ccgagcaccg ctgtttccac cggcttcctt 780  
 acttatgccg cgcacgaagg tttcaatcgt gccagctct acgaagcgat gtcacagacc 840  
 acaggcaagg gcttgtgtga tgtagcgtc ttgaggcgac ttgtcgatga agctccgaaa 900  
 40 gaaatggcgg agttgattga gacatataag gttcctgtcg ataacaccga gcgtggagt 960  
 agggcgcgcc gggcagtggg taagagcgga aaagagcttc tctccgatt ggacgcggat 1020  
 45 tacgggacgc gtggttcctt caggacacg acgggcctaa tgatggagt ctcgtcgaca 1080  
 cacggaacag cgctctatgc ccagttcgt aaagccgtga acacggcgcc aaagattcgg 1140  
 cgcgtacgcg gaagtgcgtt ggttctcgaa cccggttcca ccacggtcgg tgcgttgtc 1200  
 55 gatggcgagc cggtgacgat cgcggtcgt tcgatcatct tggcgactgg agggattcag 1260

EP 1 473 368 A1

ggcctctacg aggtcacgga taaccgcat acgctcacgg gtgatgtca tggcatggcg 1320

5 atggacgctg gcgcggagtt cgtcgacatg gagttcatgc agttctaccc gctttcagtc 1380

aatgaggagg gcgcaccgac actcttcttc tatccggatt tcccaggcg cgccaagctc 1440

10 attgacgacg ggggccgaaa cgtcctggtg aagcatctcg gcgagggtc gcaatacctt 1500

tcggagttgc ataattggga tcagctagcc gcggtggtac agacggagat tgtcgaaggc 1560

aggaaggat ttgtcgactt ccgcgaaacg aaaccggagg aatggcgcc tgattcactc 1620

20 acaggcacct tcttgggcaa gtgtgtcccg aatttcatga ccacccgggt acaggttgcg 1680

ccgtcatcgc actacacgat cggcggctcg aaagtcgat tggatggccg taccaatctt 1740

25 ccgaagctct acgctgtcgg cgagttggcc ggtggcgtgc atggcgcaa ccgccacggt 1800

ggcacggcgc tggtcgatgc catgacgtac ggccggattg ctggacggca cgcggcggga 1860

30 agcctcaacg gcgcggctgc gacgggaggt gcagcgcttc taccgcagg cagcaaagcg 1920

35 ggaaaggcaa gccggattga gggcgcaatg agcgatctgc gccgcgcaa ccagctcgct 1980

cttggtccta ttcgtgatgc cgtacggctt caacgcgttg gggagctgtt tgctgaactc 2040

40 ttggacgagg tccgtctggt cggttggaac ggctacaagg aaatgcagga aatcttgcg 2100

gtcgagcgtg cgatcaagct gtctgacgt atgcgccagg ccatgttacg ccgcacggag 2160

acacgcggag tccactatcg ggccgatttc ccgagctcca gtgatgcatg gttgaagaag 2220

50 caggatattg cattgcgcga tggggcgttg cattcaaag acgttcccct ctaatcaaag 2280

tcgcataagc gcgatattgc atcaggatatt tgtctcgtg tgggtgcatt tgtgcctcgg 2340

55

EP 1 473 368 A1

cgcccaaggg tgagagcgga aggtggagct cgccgccgcg ggcagatctg gtgtgtgtcg 2400

5 ttgttccgta tcggtagcaa aaacaatctg acttcgctag cgggcaggta agccgacggc 2460

gtatgctcgc cgaaggttct tgaattgagt tcgtggtgac catgtcgtg cgggtgtgaat 2520

10 cgctatttag gagactttat catggtaatg gcagcggtaa ttcataagaa ggggtggaccc 2580

gataacttcg tatgggagga agttaaggtt ggctcgcccg gcccggttca agtcgcactg 2640

15 cgcaatacgg ccatcggggt aaacttcctg gatacttacc accgcgcagg cattcctcac 2700

cgctggtcgc ttggcgagcc gccgattgtg gtccgcttcg aagccgccgc tgtggttgag 2760

20 gaagtcggtc ccggtgtaac cgacttcacc gttggtgagc gggctcgcac ttgtcttcgc 2820

25 cccctcgggtg cctacagcca ggagcgcctc taccgccgcg aaaaactgat caaggttcca 2880

aaggacctgg atcttgatga cgtgcacctc gcgggattga tgctcaaggc gatgacagca 2940

30 caatatctgc tgcacagac gcacaaggta aagccgggtg actacgtgtt gattcacgcg 3000

gcggctggcg gcatgggcca catcatggtt ccttgggcgc gccacctcgg cgtacogtg 3060

atcgggacgg tcagcacgga agaaaaggct gagactgtc gcaaaactcg ctgccaccat 3120

40 accatcaatt attccactca ggatttcgct gaggtagttc gcgaaatcac gggcggaag 3180

gggtgcgacg tggctacga ttccatcggg aaagacacac tccagaagtc gtcgactgt 3240

45 ctgcggccgc gcggtatgtg tgcggcctac gggcacgcat ccggcgtggc agatccgatc 3300

agggtcgtcg aggacctcgg tgtacgtgga tcgtgttca ttactagacc cgcactctgg 3360

cattacatgt cgaaccgcag tgagattgac gaagggtcga agtcctgtt cgatgccgtc 3420

55 aaggcgggtg tactccatag cagtgtcgca aagaccttc cttcgggga ggcagcggcg 3480

EP 1 473 368 A1

gcgcacaaat acatgggtgg tcgtcagacg atcggctcga ttgttttgct tccgcaagcg 3540  
5 taggtagccg tagggcgta ccccggaatt tcgggggtgac cgaaaacgtc gccgtgaacg 3600  
cctatctgcg gatgttggcg gttgggtggc attatttttc ggcagcccggt cgttcgctgc 3660  
10 cggctcgtac gttgccggtt tggacagttc gtcaacgggg cgattgtgat ttccaacccg 3720  
cgaccctcga ttgggtcaac gtgtgtttga ctttctgtga actctgtcta agtacacgag 3780  
15 aggtgtgtag atcagacgcg tgccgaagct actcggatgt tccctgtcga ggcaattgaa 3840  
agagtcaata cacatgaacg tattcaatag agagatagtg ctttccgtcc ggcactggac 3900  
20 ggataaactt ttcagcttta ccgcaacccg taacgcgcg 3938

<210> 20  
<211> 333  
30 <212> PRT  
<213> Burkholderia sp.

35 <400> 20

Met Val Met Ala Ala Val Ile His Lys Lys Gly Gly Pro Asp Asn Phe  
1 5 10 15

40 Val Trp Glu Glu Val Lys Val Gly Ser Pro Gly Pro Gly Gln Val Arg  
20 25 30

45 Leu Arg Asn Thr Ala Ile Gly Val Asn Phe Leu Asp Thr Tyr His Arg  
35 40 45

50 Ala Gly Ile Pro His Pro Leu Val Val Gly Glu Pro Pro Ile Val Val  
50 55 60

55 Gly Phe Glu Ala Ala Ala Val Val Glu Glu Val Gly Pro Gly Val Thr

EP 1 473 368 A1

	65	70	75	80
5	Asp Phe Thr Val Gly Glu Arg Val Cys Thr Cys Leu Pro Pro Leu Gly			
	85	90	95	
10	Ala Tyr Ser Gln Glu Arg Leu Tyr Pro Ala Glu Lys Leu Ile Lys Val			
	100	105	110	
15	Pro Lys Asp Leu Asp Leu Asp Asp Val His Leu Ala Gly Leu Met Leu			
	115	120	125	
20	Lys Gly Met Thr Ala Gln Tyr Leu Leu His Gln Thr His Lys Val Lys			
	130	135	140	
25	Pro Gly Asp Tyr Val Leu Ile His Ala Ala Ala Gly Gly Met Gly His			
	145	150	155	160
30	Ile Met Val Pro Trp Ala Arg His Leu Gly Ala Thr Val Ile Gly Thr			
	165	170	175	
35	Val Ser Thr Glu Glu Lys Ala Glu Thr Ala Arg Lys Leu Gly Cys His			
	180	185	190	
40	His Thr Ile Asn Tyr Ser Thr Gln Asp Phe Ala Glu Val Val Arg Glu			
	195	200	205	
45	Ile Thr Gly Gly Lys Gly Val Asp Val Val Tyr Asp Ser Ile Gly Lys			
	210	215	220	
50	Asp Thr Leu Gln Lys Ser Leu Asp Cys Leu Arg Pro Arg Gly Met Cys			
	225	230	235	240
55	Ala Ala Tyr Gly His Ala Ser Gly Val Ala Asp Pro Ile Arg Val Val			
	245	250	255	
60	Glu Asp Leu Gly Val Arg Gly Ser Leu Phe Ile Thr Arg Pro Ala Leu			
	260	265	270	



EP 1 473 368 A1

5 Trp His Tyr Met Ser Asn Arg Ser Glu Ile Asp Glu Gly Ser Lys Cys  
 275 280 285  
 Leu Phe Asp Ala Val Lys Ala Gly Val Leu His Ser Ser Val Ala Lys  
 290 295 300  
 10 Thr Phe Pro Leu Arg Glu Ala Ala Ala Ala His Lys Tyr Met Gly Gly  
 305 310 315 320  
 15 Arg Gln Thr Ile Gly Ser Ile Val Leu Leu Pro Gln Ala  
 325 330  
 20 <210> 21  
 <211> 547  
 <212> PRT  
 25 <213> Burkholderia sp.  
 <400> 21  
 30 Met Ser Asp Val Leu Val Thr Asp Val Leu Val Val Gly Glu Gly Cys  
 1 5 10 15  
 35 Ala Gly Gln Thr Ala Ala Leu Thr Ala Ser Glu Ser Gly Cys Asp Val  
 20 25 30  
 40 Ile Met Leu Gly Asp Gly Arg Ala Pro Ser Thr Ala Val Ser Thr Gly  
 35 40 45  
 45 Phe Leu Thr Tyr Ala Ala His Glu Gly Phe Asn Arg Ala Gln Leu Tyr  
 50 55 60  
 50 Glu Ala Met Ser Gln Thr Thr Gly Lys Gly Leu Cys Asp Val Ala Leu  
 65 70 75 80  
 55 Leu Arg Arg Leu Val Asp Glu Ala Pro Lys Glu Met Ala Glu Leu Ile  
 85 90 95

EP 1 473 368 A1

5	Glu Thr Tyr Lys Val Pro Val Asp Asn Thr Glu Arg Gly Val Arg Ala	100	105	110
10	Arg Arg Ala Val Gly Lys Ser Gly Lys Glu Leu Leu Ser Gly Leu Asp	115	120	125
15	Ala Asp Tyr Gly Thr Arg Gly Ser Leu Glu Asp Thr Thr Gly Leu Met	130	135	140
20	Met Glu Phe Ser Ser Thr His Gly Thr Ala Leu Tyr Ala Gln Leu Arg	145	150	155 160
25	Lys Ala Val Asn Thr Ala Pro Lys Ile Arg Arg Val Arg Gly Ser Ala	165	170	175
30	Leu Val Leu Glu Pro Gly Ser Thr Thr Val Gly Ala Leu Val Asp Gly	180	185	190
35	Glu Pro Val Thr Ile Ala Ala Arg Ser Ile Ile Leu Ala Thr Gly Gly	195	200	205
40	Ile Gln Gly Leu Tyr Glu Val Thr Asp Asn Pro His Thr Leu Thr Gly	210	215	220
45	Asp Gly His Gly Met Ala Met Asp Ala Gly Ala Glu Phe Val Asp Met	225	230	235 240
50	Glu Phe Met Gln Phe Tyr Pro Leu Ser Val Asn Glu Glu Gly Ala Pro	245	250	255
55	Thr Leu Phe Phe Tyr Pro Asp Phe Pro Arg Arg Ala Lys Leu Ile Asp	260	265	270
	Asp Gly Gly Arg Asn Val Leu Val Lys His Leu Gly Glu Gly Ser Gln	275	280	285

EP 1 473 368 A1

Tyr Leu Ser Glu Leu His Asn Trp Asp Gln Leu Ala Ala Val Val Gln  
 290 295 300  
 5  
 Thr Glu Ile Val Glu Gly Arg Lys Val Phe Val Asp Phe Arg Glu Thr  
 305 310 315 320  
 10  
 Lys Pro Glu Glu Trp Ala Pro Asp Ser Leu Thr Gly Thr Phe Leu Gly  
 325 330 335  
 15  
 Lys Cys Val Pro Asn Phe Met Thr Thr Pro Val Gln Val Ala Pro Ser  
 340 345 350  
 20  
 Ser His Tyr Thr Ile Gly Gly Leu Lys Val Asp Val Asp Gly Arg Thr  
 355 360 365  
 25  
 Asn Leu Pro Lys Leu Tyr Ala Val Gly Glu Leu Ala Gly Gly Val His  
 370 375 380  
 30  
 Gly Ala Asn Arg His Gly Gly Thr Ala Leu Val Asp Ala Met Thr Tyr  
 385 390 395 400  
 Gly Arg Ile Ala Gly Arg His Ala Ala Gly Ser Leu Asn Gly Ala Ala  
 405 410 415  
 35  
 Ala Thr Gly Gly Ala Ala Leu Leu Pro Ala Gly Ser Lys Ala Gly Lys  
 420 425 430  
 40  
 Ala Ser Arg Ile Glu Gly Ala Met Ser Asp Leu Arg Arg Ala Asn Gln  
 435 440 445  
 45  
 Leu Ala Leu Gly Pro Ile Arg Asp Ala Val Arg Leu Gln Arg Val Gly  
 450 455 460  
 50  
 Glu Leu Phe Ala Glu Leu Leu Asp Glu Val Arg Ser Phe Gly Trp Asn  
 465 470 475 480  
 55  
 Gly Tyr Lys Glu Met Gln Glu Ile Leu Arg Val Glu Arg Ala Ile Lys

EP 1 473 368 A1

485 490 495  
 5 Leu Ser Asp Ala Met Arg Gln Ala Met Leu Arg Arg Thr Glu Thr Arg  
 500 505 510

10 Gly Val His Tyr Arg Ala Asp Phe Pro Ser Ser Ser Asp Ala Trp Leu  
 515 520 525

15 Lys Lys Gln Val Phe Ala Leu Arg Asp Gly Ala Leu His Phe Lys Asp  
 530 535 540

20 Val Pro Leu  
 545

25 <210> 22  
 <211> 24  
 <212> DNA  
 <213> Artificial

30 <220>  
 <223> Forward primer for Asymmetric reductase coding region

35 <400> 22  
 caccatgtcg gatgttcttg taac 24

40 <210> 23  
 <211> 19  
 <212> DNA  
 45 <213> Artificial

<220>  
 50 <223> Reverse primer for Asymmetric reductase coding region


<400> 23  
 55 ctacgcttgc ggaagcaaa 19

## Claims


1. A gene including: DNA having a base sequence represented by SEQ ID NO: 19 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.
2. A gene including: DNA having a base sequence represented by SEQ ID NO: 17 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.
3. A gene including: DNA having a base sequence represented by SEQ ID NO: 18 that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound; or DNA that hybridizes with the DNA under stringent conditions.
4. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, **characterized by** including a DNA sequence encoding an amino acid sequence represented by SEQ ID NO: 20 and an amino acid sequence represented by SEQ ID NO: 21.
5. A gene that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising an amino acid sequence represented by SEQ ID NO: 20 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids.
6. A gene that encodes a protein having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising an amino acid sequence represented by SEQ ID NO: 21 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids.
7. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 6, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is derived from at least one microorganism selected from the group consisting of the genus Acetobacter, Actinomyces, Acinetobacter, Agrobacterium, Aeromonas, Alcaligenes, Arthrobacter, Azotobacter, Bacillus, Brevibacterium, Burkholderia, Cellulomonas, Corynebacterium, Enterobacter, Enterococcus, Escherichia, Flavobacterium, Gluconobacter, Halobacterium, Halococcus, Klebsiella, Lactobacillus, Microbacterium, Micrococcus, Micropolyspora, Mycobacterium, Nocardia, Pseudomonas, Pseudonocardia, Rhodococcus, Rhodobacter, Serratia, Staphylococcus, Streptococcus, Streptomyces, and Xanthomonas.
8. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 7, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is derived from a Pseudomonas microorganism.
9. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 7, in which the reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is originated from a Burkholderia microorganism.
10. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD810 strain.
11. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD811 strain.
12. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 8, in which the Pseudomonas microorganism is Pseudomonas sp. SD812 strain.
13. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 9, in which the Burkholderia microorganism is Burkholderia sp. SD816 strain.
14. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 13, in which the reductase has a catalytic activity to reduce a carbon-carbon double bond to produce an S-form compound chiral at an  $\alpha$ -position.

15. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 14, in which:

the  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is a compound represented by the following general formula (1)




wherein  $\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  each independently represent a hydrogen atom, a halogen atom, a linear or branched aliphatic hydrocarbon group having 1 to 6 carbon atoms, a linear or branched alkoxy group having 1 to 6 carbon atoms, a hydroxyl group, a carboxyl group, an aromatic group or a nitrogen-, oxygen-, or sulfur-containing heterocyclic group which may be substituted, and  $\text{R}^4$  represents a hydroxyl group, a linear or branched alkoxy group having 1 to 3 carbon atoms, or a primary, secondary, or tertiary amino group, provided that  $\text{R}^3$  is not a hydrogen atom; and  
a reduced compound is a compound represented by the following general formula (2)



wherein  $\text{R}^1$  to  $\text{R}^4$  have the same meanings as those defined above.


16. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 15, in which:

the  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound is an  $\alpha$ -haloacrylic acid represented by the following general formula (1)



wherein  $\text{R}^1$  and  $\text{R}^2$  represent hydrogen atoms,  $\text{R}^3$  represents a halogen atom, and  $\text{R}^4$  represents a hydroxyl

group; and  
the reduced compound is an  $\alpha$ -halopropionic acid having an S absolute configuration represented by the following general formula (2)



wherein R<sup>1</sup> to R<sup>4</sup> have the same meanings as those defined above.

17. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 16, in which R<sup>3</sup> represents a bromine atom.

18. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 16, in which R<sup>3</sup> represents a chlorine atom.

19. A reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to claim 16, in which R<sup>3</sup> represents a fluorine atom.

20. A plasmid, **characterized by** containing a DNA of a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 19.

21. A plasmid, **characterized by** containing a reductase gene for an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 19 and a gene for an enzyme functioning with an NADPH as a co-enzyme.

22. A transformant transformed with a plasmid according to claim 20 or 21.

23. A transformant including a product transformed by a plasmid according to claim 20, and a plasmid containing a gene for an enzyme functioning with an NADPH as a co-enzyme.

24. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound wherein the protein is an expression product of a reductase gene for the  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound according to any one of claims 1 to 19, or a protein having deletion, substitution, or addition of one or more amino acids thereof and having a reduction activity to an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound.

25. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound containing an amino acid sequence represented by SEQ ID NO: 20 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids in the said amino acid sequence.

26. A protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound containing an amino acid sequence represented by SEQ ID NO: 21 or an amino acid sequence having deletion, substitution, or addition of one or more amino acids in the said amino acid sequence.

27. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 631 and a base sequence selected from base sequences located downstream of a base at position 3,543 in the base sequence represented by SEQ ID NO: 19, where both the base sequences extend in opposite directions to each other.

28. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated

carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 631 and a base sequence selected from base sequences located downstream of a base at position 2,274 in base sequences represented by SEQ ID NO: 19, where both base sequences extend in opposite directions to each other.

5

29. A method of producing a gene that encodes a protein having an activity to reduce an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising using a pair of primers prepared by combining a base sequence selected from base sequences located upstream of a base at position 2,547 and a base sequence selected from base sequences located downstream of a base at position 3,543 in base sequences represented by SEQ ID NO: 19, where both base sequences extend in opposite directions so as to be reversed strands with respect to each other.

10

30. A method of reducing an  $\alpha$ -substituted- $\alpha,\beta$ -unsaturated carbonyl compound, comprising:

using a culture and/or treated product of a transformant according to claim 22 or 23.

15

20

25

30

35

40

45

50

55



Fig.1




Fig.2




Fig.3




Fig.4





Fig.5



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/01240

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl. <sup>7</sup> C12N15/53 C12N9/02, C12N5/10, C12N1/15, C12N1/19, C12N1/21, C12P7/40  According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl. <sup>7</sup> C12N15/53 C12N9/02, C12N5/10, C12N1/15, C12N1/19, C12N1/21, C12P7/40  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) JSTPlus (JOIS), SwissProt/PIR/GeneSeq, Genbank/EMBL/DDBJ/GeneSeq		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 2000-106891 A1 (Showa Denko Kabushiki Kaisha), 18 April, 2000 (18.04.00), Full text & EP 979877 A2 & US 6379935 B1	1-30
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 11 April, 2003. (11.04.03)		Date of mailing of the international search report 30 April, 2003 (30.04.03)
Name and mailing address of the ISA/ Japanese Patent Office  Facsimile No.		Authorized officer  Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)